

POLSKA AKADEMIA NAUK
ZAKŁAD ZOLOGII SYSTEMATYCZNEJ

A C T A Z O O L O G I C A
C R A C O V I E N S I A

Tom VIII

Kraków, 20 IV 1963

Nr 3

Stanisław BŁESZYŃSKI

Studies on the *Crambidae* (*Lepidoptera*). Part 40. A Review of the Genera of the Family *Crambidae* with Data on their Synonymy and Types

Materiały do znajomości *Crambidae* (*Lepidoptera*). Część 40. Przegląd rodzajów rodziny *Crambidae*, ich synonimy i typy

Материалы к познанию *Crambidae* (*Lepidoptera*). Часть 40. Просмотр родов семейства *Crambidae*, их синонимы и типы

***Achilo* AMSEL**

Chilopsis AMSEL, 1956, Bol. Ent. Venezolana, **10** (1/2): 32 (praeocc.)
Achilo AMSEL, 1957, Bol. Ent. Venezolana, **10** (3/4): (2).

Erected for a single species, namely, *Chilopsis lignella* AMSEL, 1956: 33, from Venezuela. AMSEL, in 1957, gave for this genus a new name *Achilo* AMSEL as the name *Chilopsis* AMSEL is preoccupied by *Chilopsis* Hmps., 1919.

***Acigona* HBN.**

Acigona HÜBNER, 1825, Verz. Bek. Schmett.: 342
Chiloïdes AMSEL, 1949, Bull. Soc. Fouad Ier Ent., **33**: 281.

The genus *Acigona* HBN. was erected for two species, namely, *Tinea manto* CRAMER and *Tinea cicatricella* HBN. The latter I designate as the type species of the genus *Acigona* HBN. The genus *Chiloïdes* AMSEL was erected for two species, namely, *Tinea cicatricella* HBN. and *Chilo hederalis* AMSEL. AMSEL did not designate the generic type of his *Chiloïdes* AMSEL. I designate the species *Tinea cicatricella* HÜBNER, 1924, Samml. eur. Schmett. f. 455, Europe, as the type species of the genus *Chiloïdes* AMSEL. The name *Chiloïdes* AMSEL is,

however, praeoccupied by another *Chiloïdes* BUTLER, 1881, Ann. Mag. nat. Hist., (5) 7: 392, belonging to Tortricidae. *Chiloïdes hederalis* (AMSEL) is a synonym of *Cephis galleriellus* RAG.

Adelpherupa HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 3: 541. This genus was described in Crambidae but it is referable to Schoenobiidae (BŁESZYŃSKI, 1961 d: 3).

Agriphila HBN.

Chilo ZINCKEN, 1817, Mag. Ent., 2: 34 (in part)

Agriphila HÜBNER, [1825], Verz. bek. Schmett.: 365

Pediasia HÜBNER, [1825], Verz. bek. Schmett.: 365 (in part)

Exoria HÜBNER, [1825], Verz. bek. Schmett.: 367 (in part).

The genus *Agriphila* HBN. was established for five species, of these three, *A. selasella* (HBN.), *A. deliella* (HBN.) and *A. tristella* (D. & SCHIFF.), are members of *Agriphila*. *A. deliella* (D. & SCHIFF.) was designated by LHOMME (1935: 62) as the type species of the genus *Agriphila* HBN. In the genus *Pediasia* HBN., HÜBNER placed two members of *Agriphila* HBN., *A. inquinatella* (D. & SCHIFF.) and *A. geniculea* (HAW.), and in the genus *Exoria* HBN. was placed *A. culmella* (L.).

Alamogordia DYAR & HEINRICH

Alamogordia DYAR & HEINRICH, 1927, Proc. U. S. nat. Mus., 71 (19) :38.

This genus was erected for a single species, *Diatraea parallelæ* KEARFOTT, 1908, Proc. U. S. nat. Mus., 35: 391, from New Mexico.

Amselia BŁESZ.

Amselia BŁESZYŃSKI, 1959, Beitr. naturforsch. Forsch. Südwestdeutschl., 20: 278.

Established for a single species, *Crambus heringi* AMSEL, 1935, Mitt. zool. Mus. Berl., 20: 278, from Palestine.

Anaclastis TURNER

Anaclastis TURNER, 1904, Proc. roy. Soc. Queensland, 18: 164.

Established for a single species, *Crambus apicistrigellus* MEYR., 1879, Proc. Linn. Soc. N. S. W. 4: 209, Australia. This species was placed by HAMPSON (1896: 963) in the genus *Mesolia* RAG.

***Ancalidia* J. JOAN.**

Ancalidia J. JOANNIS, 1932, Soc. ent. Fr. Livre centen. :446.

This genus was erected for one species, namely *A. nivae* J. JOAN., 1932, soc. ent. Fr. Livre centen, : 446, from Mauritius.

***Ancylolomia* HBN.**

Chilo ZINCKEN, 1817, Mag. Ent. 2: 34 (in part)

Ancylolomia HÜBNER, [1825], Verz. bek. Schmett.: 363

Jartheza WALKER, 1863, List. Spec., 27: 183

Ctenus MABILLE, 1906, Ann. Soc. ent. Fr., 75: 32

Pseudoctenella STRAND, 1907, Bull. Soc. ent. Fr., 1907: 175

Tollia AMSEL, 1949, Bull. Soc. Fouad. 1 er. Ent., 33: 280.

The genus *Ancylolomia* HBN. was erected for three species, namely, *A. disparella* (HBN.), *A. tentaculella* (HBN.), and *A. palpella* (D. & SCHIFF.). All these three species are the actual members of this genus. MOORE, in 1866: 381, designated *Ancylolomia palpella* (D. & SCHIFF.), as the type species of the genus. HAMPSON (1896: 966) designated *A. tentaculella* (HBN.) as the generic type of *Ancylolomia* HBN., but his decision is invalid. The genus *Jartheza* WALK. comprised four species, namely, *J. chrysographella* (KOLL.), *J. responsella* WALK., *J. xylinella* WALK. and *J. sabulifera* WALK. Of these, *J. responsella* WALK. is considered as a synonym of *J. xylinella* WALK., and *J. sabulifera* WALK. is sunk to the synonyms of *Chilo plejadellus* ZCK. SHIBUYA, in 1928: 145, designated *Jartheza chrysographella* (KOLL.) as the generic type of *Jartheza* WALK. HAMPSON (1896: 966) sank *Jartheza* WALK. to the synonyms of *Ancylolomia* HBN. and his opinion is right as both *A. palpella* (D. & SCHIFF.) and *A. chrysographella* (KOLL.) seem to be congeneric. The genus *Ctenus* MAB. was erected for one species, viz. *C. malacellus* MAB., 1906, Ann. Soc. ent. Fr. 74: 32, from Malaga. STRAND gave a nomen novum *Pseudoctenella* STRAND as the name *Ctenus* MAB. is preoccupied by *Ctenus* WALCKENAER, 1905, Arachnida. *Ctenus malacellus* MAB. is a synonym of *Ancylolomia inornata* STGR., which is congeneric with *A. palpella* (D. & SCHIFF.), in consequence, MABILLE's genus falls into synonyms of *Ancylolomia* HBN. (SHIBUYA, 1928: 145). *Tollia* AMSEL was erected for *Ancylolomia pectinatella* (ZELL.), 1847: 747 South Europe. The present writer sunk the AMSEL genus under *Ancylolomia* HBN. (1962 d: 32).

***Angustalius* MARION**

Angustalius MARION, 1954, Mém. Inst. Sci. Madagascar, (E) 5: 50.

Established for one species, viz., *A. ditaeniellus* MARION, 1954: 50, from Madagascar.

Anomocrambus BŁESZ.

Anomocrambus BŁESZYŃSKI, 1961, Pol. Pis. ent., **31**: 4.

This genus was erected for a single species, *A. homerus* BŁESZ. 1961 b: 204, from Natal.

Aphrophantis MEYR.

Aphrophantis MEYRICK, 1933, Exot. Micr., **4**: 379.

Erected for a single species, namely *A. velifera* MEYR., 1933: 379, from Fiji.

Araxates RAG. see *Talis* GUEN.

Araxes STEPH. see *Euchromius* GN., *Platytes* GN.

Araxes WALKER see *Culladia* MOORE

Archigalleria RTHSCH. see *Cephis* RAG.

Arequipa WALK. see *Crambus* F.

Argentochiloïdes BŁESZ.

Argentochiloïdes BŁESZYŃSKI, 1961, Zeit. Wien. ent. Ges., **46**: 36.

This genus was erected for a single species, namely *A. xanthodorsellus* BŁESZ. 1961c: 36, from Tanganyika-Territory.

Argyria HBN.

Argyria HÜBNER, [1825], Verz. bek. Schmett. :372

Urola WALKER, 1863, List Spec., **27**: 181.

This genus was erected for two species, *A. pusillalis* (HBN.), 1818, Zutr., figs. 167, 168, and *A. nummulalis* (HBN.), 1818, l.c. figs. 185, 186. The former is a synonym of *A. lacteella* (F.) and the latter is a synonym of *A. argentana* (MARTYN). FERNALD in 1896: 70 designated *Argyria nummulalis* (HBN.) as the type species of *Argyria* HBN. The genus *Urola* WALK. comprised six species, of which five are referable to *Argyria* HBN. I designate *Urola michrochysella* [sic.] WALK., 1863: 181, as the type species of *Urola* WALK. Thus, *Urola* WALK. falls to the synonyms of *Argyria* HBN. HAMPSON in 1896: 943 combined a cluster of five genera (*Platytes* GUEN., *Argyria* HBN., *Catharylla* ZELL., *Urola* WALK., *Ptychopseustis* MEYR.) into one genus *Platytes* GUEN., but his decision is obviously wrong. The genus *Argyria* HBN. is perfectly distinct from *Platytes* GUEN. The latter appears to be distributed only in the Pala-

earctic Region, and the most part of the members of the former are indigenous to the Nearctic and Neotropical Regions. In fact, HAMPSON made an extraordinary chaos in the group in question, and all the genera placed by him under *Platytes* GUEN. are in need of thorough revision.

***Aurelianus* BŁESZ.**

Aurelianus BŁESZYŃSKI, 1961, Coridon (A) 3: 2.

Aurelianus BŁESZ. was erected for two species, namely, *Chilo discellus* WALK., 1863: 141, from South Africa and *Crambus mesoscia* Hmps., 1919: 279, also from South Africa. *Chilo discellus* WALK. has hitherto been considered to be a member of the genus *Crambus* F. Type species, *Chilo discellus* WALK., by original designation.

***Aureocramboides* BŁESZ.**

Aureocramboides BŁESZYŃSKI, 1961, Pol. Pis. ent. 31: 205.

This genus was erected for a single species, namely, *A. apollo* BŁESZ. 1961b: 205, from British East Africa.

***Autarotis* MEYR.**

Autarotis MEYRICK, 1886, Trans. ent. Soc. Lond., 1886: 269.

Described for a single species, namely *A. euryala* MEYR., 1886: 270, from Fiji.

***Bissetia* KAPUR**

Bissetia KAPUR, 1950, Trans. R. ent. Soc. Lond., 101: 417.

This genus was erected for a single species, *Chilo steniellus* HAMPSON, 1899: 305, India.

***Bleszynskia* DE LATTIN**

Crambopsis DE LATTIN, 1952, Ent. Zeit., 62: 91 (praeoc.)
Bleszynskia DE LATTIN, 1961, Ent. Zeit., 71: 115.

Erected for one species, namely, *Crambus malacellus* DUP. The name *Crambopsis* DE LATT. is preoccupied by *Crambopsis* WALK., 1865: 634, in Noctuidae.

Caffrocrambus BŁESZ.

Caffrocrambus BŁESZYŃSKI, 1961, Pol. Pis. ent., 31: 200.

Erected for three species, namely, *Crambus chalcimerus* HMPS., 1919: 285, Basutoland, and *C. dichotomellus* HMPS., 1919: 284, South Africa and *Caffrocrambus alcibiades* BŁESZ., South Africa. Type species, *Crambus dichotomellus* HMPS., by original designation.

Calamotropha ZELL

Calamotropha ZELLER, 1863, Chil. Cramb. Gen. Spec. :8

Myzea WALKER, 1863, List. Spec., 27: 190

Conocrambus HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 4: 443 (in part).

This genus was erected for two species, namely *Tinea paludella* HBN., 1824 and *Chilo aureliellus* F. R., 1834: 261, Europe. Of these, *Calamotropha paludella* (HBN.) was designated by LHOMME, 1935: 577, as the type species of *Calamotropha* ZELL. The genus *Myzea* WALK. was described for one species, *M. tonsalis* WALK., 1863: 190, from Borneo. *Myzea* WALK. was synonymized with *Calamotropha* ZELL. by the writer in 1961a: 151. *Conocrambus* HMPS. was erected for three species, *C. artemietellus* HMPS., New Guinea, *C. calamosus* HMPS., Transvaal, and *C. obliqua* HMPS., Brazil. All these three species are apparently members of three genera. Of these, *C. calamosus* HMPS. is a synonym of *Calamotropha paludella* (HBN.) (BŁESZYŃSKI, 1961a: 151). For *C. obliqua* HMPS. was erected the genus *Vaxi* BŁESZ., 1962.

Calarina WALK. see *Surratha* WALK.

Canuza WALKER, 1866, List Spec., 35: 1771 (= *Erotomanes* MEYR. 1882, Proc. Linn. Soc. N. S. Wales, 7: 152). Described in *Crambidae* but referable to *Pyraustidae* (BŁESZYŃSKI, 1962 a: 3).

Carvanca WALK., see *Pediasia* HBN.

Catancyla HMPS., see *Diadexia* TURN.

Catharylla ZELL.

Catharylla ZELLER, 1863, Chil. Cramb. Gen. Spec.: 50

This genus is still confused. MEYRICK in 1879: 215 sank *Catharylla* ZELL. under *Argyria* HBN. However, such a decision may be considered as invalid as MEYRICK did not designate the type species of *Catharylla* ZELL. and this genus comprised nine species of at least three distinct genera. Only in 1924: 131, SCHAUß designated *Crambus tenellus* ZELL., 1839: 174, from Brazil, as the type species of the genus *Catharylla* ZELL. This designation is valid as

Crambus tenellus ZELL. was placed by ZELLER in his genus. The remaining species of *Catharylla* ZELL. need verification. Some of those belong to the genus *Argyria* HBN.

***Catoptria* HBN.**

Tinea LINNAEUS, 1758, Syst. Nat. (ed. 10), 1: 534 (in part),
Chilo ZINCKEN, 1817, Mag. Ent., 2: 34 (in part)
Catoptria HÜBNER, [1825], Verz. bek. Schmett. :364
Argyroteuchia HÜBNER, [1825], Verz. bek. Schmett. :364
Exoria HÜBNER, [1825], Verz. bek. Schmett., :367 (in part).

ZINCKEN, in his *Chilo* ZCK. placed several actual members of the genus *Catoptria* ZCK. of these *C. verella* ZCK. was described as new to science. FABRICIUS, in his *Crambus* F., placed 52 species of which several are actual members of the genus *Catoptria* HBN. In the genus *Exoria* HBN. were placed three species, namely *Tinea combinella* D. & SCHIFF., *T. convolutella* D. & SCHIFF. and *Tinea culmella* L. I designate *Tinea combinella* D. & SCHIFF., as the type species of the genus *Exoria* HBN. This species is congeneric with *Catoptria speculalis* HBN. [1825]: 365, which is the type species of the genus *Catoptria* HBN. (designated by the present writer, in 1956: 213). In 1957: 306, the present writer cited erroneously *Crambus permuatellus* H.-S., 1848: 64, as the generic type of the genus under consideration.

***Cephis* RAGONOT**

Cephis RAGONOT, 1892, Dtsch. ent. Zeit. Lep., 5: 295
Stenochilo HAMPSON, 1896, Proc. zool. Soc. Lond., 1895: 950
Chilooides AMSEL, 1949, Bull. Soc. Fouad. ler Ent., 33: 281 (in part)
Hombergia J. JOANNIS, 1910, Bull. Soc. ent. Fr., 1910: 270.

The genus *Cephis* RAGONOT, was erected for a single species, namely *C. gallorellus* RAG., 1892: 295, from Tunis. MARTIN, in 1956: 164 sank the genera, *Stenochilo* HMPS. and *Hombergia* J. JOAN. under *Cephis* RAG. *Stenochilo* HMPS. was erected for a single species. *S. canicostalis* HMPS., 1896: 950, Hillaya, Sind; and *Hombergia* J. JOAN. was described for one species, *H. unicolor* J. JOAN., 1910: 270, from France. *Chilooides* AMSEL., 1949: 281, was established for two species, namely *Chilo hederalis* AMSEL. and *Tinea cicatricella* HBN. The former was sunk to the synonyms of *Stenochilo canicostalis* HMPS. by the writer (1959: 115), and the latter is previously designated as the type species of the genus *Acigona* HBN.

Chaerecla WALK., see *Eschata* WALK.

Chalcoëla ZELLER, 1872, Verh. zool.-bot. Ges., 22: 528. Described in *Crambidae* but referable to *Pyraustidae* (FORBES, 1920: 215; BŁĘSZYŃSKI, 1962a: 3).

***Charitopepla* MEYR.**

Charitopepla MEYRICK, 1933, Exot. Micr., 4: 379.

Established for one species, *Charitopepla drosocapna* MEYR. 1933: 379, from Congo.

***Charltona* SWINH.**

Charltona SWINHOE, 1885, Proc. Zool. Soc. Lond., 1885: 878.

Established for one species, namely *Charltona kala* SWINH., 1885: 879, from India.

***Charltoniada* STRAND**

Charltoniada STRAND, 1919, Stett. ent. Ztg., 79 (1918): 259.

This genus was established for one species, namely *Ch. difficilis* STRAND, 1919: 259, from Formosa. SHIBUYA, in 1928: 52, sank *Ch. difficilis* STRAND under *Crambus apicenotatus* HMPS., 1919: 287 (the exact date of the STRAND publication is July, 1919, and that of the HAMPSON work is March, 1919) and he placed the species under the genus *Diatraea* HMPS. However, *Crambus apicenotatus* HMPS. is not congeneric with the genus *Diatraea* GUILD. and the genus *Charltoniada* STRAND is valid.

***Chilo* ZCK.**

Chilo ZINCKEN, 1817, Mag. Ent., 2: 34

Diphryx GROTE, 1882, Bull. Z. S. Geol. Surv., 6: 273

Silveria DYAR, 1925, Insec. inscit., 13: 10

Chilotraea KAPUR, 1950, Trans. R. ent. Soc. Lond., 101: 402.

The genus *Chilo* ZCK. was erected for a cluster of species belonging actually to several genera. Of these, only one, viz. *phragmitellus* (HBN.) is an actual member of the genus in question. DUPONCHEL, in 1836: 14, designated the species *Tinea phragmitella* HBN., 1805, as the type species of *Chilo* ZCK. The genus *Diphryx* GROTE was erected for a single species *Diphryx prolatella* GROTE, 1882: 273, Wisconsin. This species was sunk by FERNALD, in 1888: 45, under *Chilo plejadellus* ZCK., 1821: 251. The genus *Silveria* DYAR was described for two species, namely *Silveria adelphilia* DYAR, and *S. hexhex* DYAR, both from Mexico. Type species, *S. hexhex* DYAR, 1925: 11, by original designation. The present writer, in 1962 d: 107, sank the DYAR genus under *Chilo* ZCK. The genus *Chilotraea* KAPUR was established for four species: *Chilo infuscatellus* SNELL.,

Chilo ceylonicus HMPS., *Chilo auricilia* DUDG. and *Chilotraea bandra* KAPUR. Of these, *Chilo infuscatus* SNELL., 1890: 94 was designated by KAPUR as the type species of *Chilotraea* KAPUR. The present writer in 1962a: 1, pointed out that both *Chilo* ZCK. and *Chilotraea* KAPUR are obviously synonyms.

HAMPSON, in 1896: 954, placed under *Chilo* ZCK. two other genera, namely *Donacoscaptes* ZELL. and *Epina* WALK. *Donacoscaptes* ZELL. was described by ZELLER for a single species, *Donacoscaptes validus* ZELL., 1877: 16, from South America. However, both *Donacoscaptes validus* ZELL. and *Chilo phragmitellus* (HBN.) are obviously not congeneric, and the ZELLER genus should preserve its validity. The genus *Epina* WALK. was erected for one species, *Epina dichromella* WALK., 1866: 1707 (?North America). KAPUR, in 1948: 191, pointed out that the genus *Epina* WALK., is valid, and sank to its synonymy the genus *Diatraenopsis* DYAR & HEINRICH, 1927: 39.

The genus *Chilo* ZCK. is still in need of a thorough revision as it includes many species not congeneric with *Chilo phragmitellus* (HBN.). Most of the species from the Nearctic and Neotropical Regions described in this genus are referable to different genera. On the other hand, many Ethiopian and Oriental species described in *Diatraea* GUILD. are obviously members of the genus *Chilo* ZCK.

Chilooides AMSEL, see *Acigona* HBN.

Chilopsis HMPS., see *Myelobia* H.-S.

Chilopsis AMSEL, see *Achilo* AMSEL

Chilotraea KAPUR, see *Chilo* ZCK.

Chrysocramboides BŁESZ., see *Chrysocrambus* BŁESZ.

***Chionobosca* TURNER**

Chionobosca TURNER, 1911, Ann. Queensland Mus., 10: 114.

This genus was erected for a single species, namely *Chionobosca actinopis* TURNER, 1911: 114, Port Darwin, Australia.

***Chrysocrambus* BŁESZ.**

Crambus FABRICIUS, 1798, Ent. Syst. Suppl.: 464 (in part)

Thisanotia HÜBNER, [1825], Verz. bek. Schmett.: 366

Chrysocrambus BŁESZYŃSKI, 1957, Acta zool. crac., 1: 443

This genus was erected for five species, of which *Crambus cassentiniellus* ZELL. 1849: 312, Europe, was designated as the type species. However, the reference on the latter species was given incorrectly as the first author of *Crambus cassentiniellus* is HERRICH-SCHÄFFER, 1848: 49. The genus *Chrysocrambus* BŁESZ. was divided into two subgenera, *Chrysocrambus* BŁESZ. s. str. and *Chrysocramboides* BŁESZ., subgeneric type *Phalaena craterella* SCOP., 1763: 246.

Chrysoteuchia HÜBNER, see *Crambus* F.

Classeya BŁESZ.

Classeya BŁESZYŃSKI, 1960, Čas. česk. Spol. ent., 57: 267.

This genus was erected for six species of the genera *Argyria* HBN., *Crambus* F. and *Platytes* GUEN. Generic type *Argyria bicuspidalis* HMPS., 1919: 446, Central Africa, by original designation.

Coenotalis HMPS., see *Parerupa* HMPS.

Coniesta HMPS., see *Haimbachia* DYAR.

Conocrambus HMPS.

Conocrambus HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 3: 443.

This genus was erected for three species, namely *Conocrambus atrimictellus* HMPS., 1919: 443, New Guinea, *C. calamosus* HMPS., 1919, Transvaal, and *C. obliqua* HMPS., 1919, Brazil. Generic type, *C. atrimictellus* HMPS., by original designation. All the three species of this genus are not congeneric with each other. *Crambus calamosus* HMPS. is sunk to the synonyms of *Calamotropha paludella* (HBN.) (BŁESZYŃSKI, 1961a: 152). For *Conocrambus obliqua* HMPS. is erected the genus *Vaxi* BŁESZ.

Conotalis HMPS.

Conotalis HAMPSON, 1919. Ann. Mag. nat. Hist., (9) 4: 150.

This genus was established for one species, namely *Charltona aurantifascia* HMPS., 1896: 479, South Africa.

Corynophora BERG

Halterophora MEYRICK, 1897, Trans. ent. Soc. Lond., 1897: 378 (praeoc.).

Corynophora BERG, 1898, Com. Mus. Buenos Aires, 1: 42.

The genus *Halterophora* MEYR. was described for one species, viz. *Crambus lativittalis* WALK., 1863: 171 (= *Crambus halterellus* ZELL.), from Sydney. BERG replaced the MEYRICK name by *Corynophora* BERG as *Halterophora* MEYR. was preoccupied.

Crambidiatraea Box & CAPPS

Crambidiatraea Box & CAPPS, 1955, Proc. R. ent. Soc. Lond., (B) 24: 175.

Erected for four species, *Diatraea cayenella* DYAR & HEINR., *D. strigipennella* DYAR, *D. castrensis* DYAR & HEINR. and *D. entreriana* Box. Type species, *D. cayenella* DYAR & HEINR., 1927: 27, from French Guiana, by original designation.

Grambidion MAB., see *Culladia* MOORE

Crambopsis DE LATTIN, see *Bleszynskia* DE LATTIN

Crambus F.

Tinea LINNAEUS, 1758, Syst. Nat. (ed. 10) 1: 534 (in part)

Crambus FABRICIUS, 1798, Ent. Syst. Supp.: 464

Palparia HAWORTH, 1811, Lep. Brit., (3): 481

Chilo ZINCKEN, 1817, Germ. Mag. Ent., 2:34 (in part)

Argyroteuchia HÜBNER, [1825], Verz. bek. Schmett.: 364 (in part)

Agriphila HÜBNER, [1825] Verz. bek. Schmett.: 365 (in part)

Chrysoteuchia HÜBNER, [1825], Verz. bek. Schmett.: 366

Selagia HÜBNER, [1825], Verz. Bek. Schmett.: 367 (in part)

Arequipa WALKER, 1863, List Spec., 27: 195.

LINNAEUS in his *Tinea* placed two actual members of the genus *Crambus* F., namely *C. pascuellus* (L.) and *C. pratellus* (L.). The genus *Crambus* F. comprised 62 species, of which only few are actually species of the genus *Crambus* F. s. str. The genus *Palparia* HAW. contained also few species of the genus in question. I designate *Tinea pascuella* L. as the type species of the genus *Palparia* HAW., thus the latter falls into synonyms of the genus *Crambus* F. s. str. The genus *Chilo* ZCK. comprised some species of the genus discussed, e. g., *Crambus pascuellus* (L.), *C. pratellus* (L.) and others. The genus *Argyroteuchia* HBN. comprised nine species, of which only two are not congeneric with *Crambus pascuellus* (L.), which is the generic type of the genus *Crambus* F. s. str.; these are *Catoptria falsella* (D. & SCHIFF.) and *Platytes alpinellus* (HBN.). I designate *Tinea pascuella* L. as the type species of *Argyroteuchia* HBN. The genus *Agriphila* HBN. comprised five species, one of which was named by HÜBNER *Agriphila pratalis*. It might be considered to be *Crambus dumetellus* (HBN.), however, the matter is obscure and difficult to solve. The genus *Chrysoteuchia* HBN. was erected for two species: „*Ch. hortalis*“ and „*Ch. caricetalis*“. The two are conspecific, *Crambus hortuellus* (HBN.). The genus *Selagia* HBN. comprised three species, of these *Crambus perlellus* (SCOP.) is congeneric with *C. pascuellus* (L.). The genus *Arequipa* WALK. was erected for one species, *Arequipa turbatella* WALK., 1863: 196, North America, which is congeneric with *Crambus pascuellus* (L.). *Tinea pascuella* L. was designated as the generic type of the genus *Crambus* F. by CURTIS in 1825: 109.

Crunophila MEYR. see *Ubida* WALK.
Cryptomima MEYR. see *Gadira* WALK.
Ctenus MAB. see *Ancylolomia* HBN.

***Culladia* MOORE**

Araxes WALKER, 1863, List. Spec., 27: 192 (praeoc.)
Culladia MOORE, 1886, Lep. Ceyl., 3: 383
Crambidion MABILLE, 1899, Ann. Soc. ent. Fr., 68: 748.

The genus *Araxes* WALK. was erected for four species, namely *Araxes admigratella* WALK., 1863: 192, *A. caesella* WALK., 1863: 193, *A. celsella* WALK., 1863: 193 and *A. decursella* WALK., 1863: 194, all from Ceylon. Of these, *A. caesella* WALK. is a synonym of *Calamotropha famulella* (WALK.) (SYN. N.), *A. decursella* WALK. belongs to the subfamily Pyralinae, and *A. celsella* WALK. is unrecognized by the present writer (type lost). The name *Araxes* WALK. is preoccupied by *Araxes* STEPHENS, 1834: 315; and MOORE, in 1886, gave a new name *Culladia* MOORE, for the WALKER genus. MOORE designated the species *A. admigratella* WALK. as the generic type of the genus *Culladia* MOORE. The genus *Crambidion* MAB. was established for a single species, namely *C. achroellum* MAB., 1899: 748, from Madagascar. SHIBUYA, in 1928: 44 places *Crambidion* MAB. under *Culladia* MOORE and cites *Crambus incospicuellus* SNELL., 1872: 102, from Lower Guinea, to be the generic type of *Crambidion* MAB. Probably, SHIBUYA supposed both *Crambus incospicuellus* SNELL., and *Crambidion achroellum* MAB. to be identical species. However, such a supposition needs a confirmation based on a study of the types of the two species. On the other hand, however, *Crambidion achroellum* MAB. is obviously congeneric with *Culladia admigratella* (WALKER).

***Cypholomia* MEYR.**

Cypholomia MEYRICK, 1933, Exot. Micr., 4: 378

Established for a single species, namely, *Cypholomia leptodeta* MEYR. 1933: 378, from Congo.

***Deuterolia* DYAR**

Deuterolia DYAR, 1914, Proc. U. S. nat. Mus., 47: 402.

Erected for a single species viz. *Deuterolia nipis* DYAR, 1914: 402, from Mexico.

Diadexia TURNER

Diadexia TURNER, 1905, Proc. roy. Soc. Queensland, **19**: 56
Catancyla HAMPSON, 1919, Ann. Mag. nat. Hist., (9) **3**: 140.

Erected for a single species, viz. *Diadexia parodes* TURNER, 1905: 56, Australia. *Catancyla* HMPS. was erected for one species, *C. brunnea* HMPS., 1919: 140 from Australia. This species is a synonym of *Diadexia parodes* TURN.

Diatraea GUILD.

Diatraea GULDING, 1828, Trans. Soc. Encour. Arts, etc., **46**: 148.
Diatraerupa SCHAUS, 1913, Ann. Mag. nat. Hist., (8) **11**: 240.
Iesta DYAR, 1909, Proc. ent. Soc. Wash., **11**: 29
Trinidadia DYAR & HEINRICH, 1927, Proc. U. S. nat. Mus., **71** (19): 5.

The genus *Diatraea* GUILD. was erected for a single species, namely, *Phalaena saccharalis* F., 1794: 238. The genus *Diatraerupa* SCHAUS, was erected for one species, *Diatraerupa guapilella* SCHAUS, 1913: 240, from Costa Rica. This species was transferred by SCHAUS, 1922: 137, to the genus *Iesta* DYAR. The latter was erected for one species, *Iesta lisetta* DYAR, 1909: 29, from Florida. The genus *Trinidadia* DYAR & HEINR. was established for one species, namely *Diatraea minimifacta* DYAR, 1911: 202, SOUTH America. The genera *Diatraerupa* SCHAUS, *Iesta* DYAR, and *Trinidadia* DYAR & HEINR. were sunk to the synonyms of the genus *Diatraea* GUILD., by Box, in 1931: 11.

Diatraenopsis DYAR & HEINR. see *Epina* WALK.
Diatraerupa SCHAUS, see *Diatraea* GUILD.

Dicymolomia ZELLER, 1872, Verh. zool.-bot. Ges., **22**: 530. Erected for one species, namely, *D. decora* ZELL., 1872: 131 from Texas. Referable not to *Crambidae* but to *Pyraustidae* (FORBES, 1920: 215; BŁĘSZYŃSKI, 1961 a: 3).

Diphryx GROTE, see *Chilo* ZCK.

Diploschistis MEYR.

Diploschistis MEYRICK, 1937, Exot. Micr., **5**:140.

Erected for one species, namely *Diploschistis stygiocrena* MEYR., 1937: 159, from Congo.

Diplotalis HMPS.

Diplotalis HAMPSON, 1919, Ann. Mag. nat. Hist., (9) **4**: 151.

Erected for a single species, *Diplotalis metallescens* HMPS., 1919: 152, from Nigeria.

***Diptychophora* ZELL.**

Diptychophora ZELLER, 1866, Ent. Ztg. Stettin, **27**: 153
Ditomoptera HAMPSON, 1893, Ill. Het., **9**: 179.

The genus *Diptychophora* ZELL. was erected for one species, namely *Diptychophora kuhlweini* ZELL., 1866: 144, Brazil. *D. kuhlweini* ZELL. was sunk under *D. azanalis* WALK., by MEYRICK, 1931: 107. The genus *Ditomoptera* HMPS. was erected for a single species, *Ditomoptera minutalis* HMPS., 1893: 179, from Sikkim and Ceylon. This genus was sunk under the genus in question by HAMPS., in 1896: 942.

***Donacoscaptes* ZELL.**

Donacoscaptes ZELLER, 1877, Horae Soc. ent. ross., **13**: 16.

This genus was erected as a subgenus of *Chilo* ZCK. with a single species *Donacoscaptes validus* ZELL., 1877: 16, South America. HAMPS., in 1896: 954 sank this genus under *Chilo* ZCK., however, this decision was incorrect as both *Donacoscaptes validus* ZELL. and *Chilo phragmitellus* (HBN.) (type species of *Chilo* ZCK.) are not congeneric. Thus, the ZELLER genus should preserve its validity.

Doratoperas HMPS., see *Myelobia* H.-S.

***Drasa* KAPUR**

Drasa KAPUR, Trans. R. ent. Soc. Lond., **101**: 409.

Erected for a single species, *Diatraea cashmirensis* HMPS., 1919: 543, from Kashmir.

***Elethyia* RAG.**

Elethyia RAGONOT, 1888, Ann. Soc. ent. Fr., (6) **8**: 281.

Erected for a single species, *Eromene?* *subscissa* CHRIST., 1877: 277, from Krasnovodsk.

***Eodiatraea* Box**

Eodiatraea Box, 1953, Proc. R. ent. Soc. Lond., (B) 22: 178.

Erected for four species, namely *Chilo centrellus* MÖSCHL., *Diatraea ammenella* DYAR, *D. amazonica* Box, and *D. rufescens* Box. The type species is *Chilo centrellus* MÖSCHL., 1883: 360, from Paramaribo, Surinam, by original designation.

Eoreuma ELY, see *Haimbachia* DYAR

***Epichilo* RAG.**

Epichilo RAGONOT, 1888, Ann. Soc. ent. Fr., (6) 8: 275.

Erected for a single species, namely *Epichilo parvulus* RAG., 1888: 278, India.

***Epina* WALK.**

Epina WALKER, 1866, List. Spec., 35: 1707.

Diatraenopsis DYAR & HEINRICH, 1927, Proc. U. S. nat. Mus., 71 (19): 39.

The genus *Epina* WALK. was established for one species, namely *Epina dichromella* WALK., 1866: 1707, (locality not given). The genus *Diatraenopsis* DYAR & HEINR. was described for two species, namely, *Diatraea differentialis* FERN. and *D. idalis* FERN. Type species *Diatraea differentialis* FERN., 1888: 120, Florida, by original designation.

Eromene HBN., see *Euchromius* GUEN.

Erotomanes MEYR., see *Canuza* WALK.

***Erupa* WALK.**

Erupa WALKER, 1864, List. Spec., 30: 980

Gabalanca WALKER, 1866, List. Spec., 35: 1743

Zolca WALKER, 1866, List. Spec., 35: 1769.

The genus *Erupa* WALK. was described for a single species, namely, *Erupa chilooides* WALK., 1864: 980, from Brazil. The genus *Gabalanca* WALK. was erected also for one species, *Gabalanca bilineatella* WALK., 1866: 1743, from Brazil; and the genus *Zolca* WALK. was erected for *Zolca congruella* WALK., 1866: 1769, Brazil. HAMPSON, in 1896: 951, sank the genera *Gabalanca* WALK. and *Zolca* WALK. under *Erupa* WALK.

***Eschata* WALK.**

Eschata WALKER, 1856, List. Spec., 9: 133
Chaerecla WALKER, 1865, List. Spec., 32: 633.

Eschata WALK. was erected for *Eschata gelida* WALK., 1856: 133, from India; and *Chaerecla* WALK. was described for also a single species, namely, *Chaerecla chrysargyria* WALK., 1865: 634, Ceram. HAMPSON, in 1896: 961, sank *Chaerecla* WALK. under *Eschata* WALK.

***Euchromius* GUEN.**

Eromene HÜBNER, [1825], Verz. bek. Schmett.: 366 (praeoc.)
Araxes STEPHENS, 1834, Ill. Brit. Ent. Haust., 4: 315
Palparia HAWORTH, 1811, Ent. Brit. 3: 481 (in part)
Euchromius GUENÉE, 1845, Ann. Soc. ent. Fr., (2) 3: 324
Ommatopteryx KIRBY, 1897, Handbook Lep., 5: 274.

The genus *Eromene* HBN. was erected for one species, *Tinea bella* HBN., 1796, Europe. However, the name *Eromene* HBN. is preoccupied by another *Eromene* HBN., 1821: 256, in *Noctuidae*. The genus *Araxes* STEPH. comprised a heterogeneous cluster of four species belonging to two families, *Crambidae* and *Phycitidae*, including *Palparia ocellea* HAW., which is here designated as the type species of *Araxes* STEPH. (this species is congeneric with *Euchromius bellus* (HBN.). The genus *Euchromius* GUEN. was established for three species, *Tinea bella* HBN., *Crambus ramburiellus* DUP. and *Phycis funiculellus* TR. *Euchromius* GUEN. has been considered by some workers to be a homonym, as there is a genus *Euchromia* HBN., 1819, and another *Euchromia* STEPH., 1829. However, this opinion is incorrect and the name of GUENÉE should preserve its validity. Basing on the incorrect opinion, KIRBY gave a new name *Ommatopteryx* KIRBY for the supposed homonym, *Euchromius* GUEN. For a long time, the KIRBY name has been adopted by some workers. The type species of *Euchromius* GUEN. is *Tinea bella* HBN., designated by HAMPSON, in 1896: 949. Some authors considered the species *Euchromius ocelleus* (HAW.) to be the type species of this genus, but such an opinion is invalid.

Eudorina SNELL., see *Syncrotaula* MEYR.

***Eufernaldia* HULST**

Eufernaldia HULST, 1900, J. N. Y. ent. Soc., 8: 224.

This genus was established for one species, namely, *Eufernaldia argenteonervella* HULST, 1900: 224, from Arizona.

Eugrotea FERNALD, see *Mesolia* RAG.

***Euparolia* DYAR**

Euparolia DYAR, 1914, Proc. U. S. nat. Mus., 47: 402.

Erected for one species, namely, *Euparolia nipimidalis* DYAR, 1914: 402, from Mexico.

***Eurythma* TURNER**

Eurythma TURNER, 1904, Proc. roy. Soc. Queensland, 18: 168.

Erected for a single species, namely, *Platytes latifasciella* Hmps., 1919: 445, from Port Darwin, Australia.

Exoria Hbn., see *Catoptria* Hbn., *Agriphila* Hbn.

***Exsilirarcha* SALMON & BRADLEY**

Exsilirarcha SALMON & BRADLEY, 1956, Rec. Dom. Mus., 3 (1): 73.

This genus was established for a single species, namely, for *Exsilirarcha graminea* SALMON & BRADLEY, 1956: 75, from Campbell Island (Antipodes Islands).

***Fernandocrambus* AUR.**

Fernandocrambus AURIVILLIUS, 1922, Nat. Hist. of Juan Fern., (3) 2: 263.

This genus was described for three species, *Fernandocrambus bækströmi* AUR., *F. brunneus* AUR. and *F. fuscus* AUR., 1922: 264, from Masatierra, Juan Fernandez. I designate *Fernandocrambus bækströmi* AUR., as the type species of the genus in question.

***Flavocrambus* BLESZ.**

Flavocrambus BŁESZYŃSKI, 1959, Tinea, 5: 275.

Established for two species, namely, *Crambus striatellus* LEECH and *C. aridalis* SOUTH. Type species *Crambus striatellus* LEECH, 1889: 107, from Japan, by original designation.

Gabalanea WALK., see *Erupa* WALK.

Gadira WALK.

Gadira WALKER, 1866, List. Spec., 35: 1742
Cryptomima MEYRICK, 1882, Trans. N. Zeal. Inst., 1882: 8.

The genus *Gadira* WALK. was erected for a single species, namely, *Gadira acerella* WALK., 1866: 1742, from New Zealand. MEYRICK erected his *Cryptomima* MEYR. also for *Gadira acerella* WALK. HAMPSON, in 1896: 917, sank *Cryptomima* MEYR. under *Gadira* WALK.

Girdharia KAPUR

Girdharia KAPUR, 1950, Trans. R. ent. Soc. Lond., 101: 428.

Erected for a single species, namely, *Girdharia tauromma* KAPUR, 1950: 428, from India.

Haimbachia DYAR

Haimbachia DYAR, 1909, Proc. ent. Soc. Wash., 11: 28
Eoreuma ELY, 1910, Proc. ent. Soc. Wash., 12: 204
Coniesta HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 3: 540.

Haimbachia DYAR was established for one species, *Crambus placidellus* HAIMB., 1907: 44, from New Jersey. *Eoreuma* ELY was erected for *Chilo densellus* ZELL., 1881: 158, from Mexico. *Coniesta* HMPS. was described for two species, namely, *Chilo araealis* HMPS., 1912: 1250, from Madras and *Coniesta undilinealis* HMPS., 1919: 540, British Central Africa. Type species *Chilo araealis* HMPS., by original designation. *Coniesta undilinealis* HMPS. is not congeneric with *C. araealis* (HMPS.), being referable to *Schoenobiidae*. Both *Eoreuma* ELY and *Coniesta* HMPS. were sunk under *Haimbachia* DYAR by the present writer (1961d: 1). The genus *Haimbachia* DYAR is very numerous with the species and is widely distributed. Many of its species were described in such genera as *Argyria* HBN., *Chilo* ZCK., *Erupa* WALK. or *Diatraea* GUILD.

Halterophora MEYR., see *Corynophora* BERG

Hednota MEYR.

Hednota MEYRICK, Trans. ent. Soc. Lond., 1886: 270.

This genus was erected for several Australian species. The species *Crambus bifractellus* WALK., 1863: 174, from South Australia was cited by Zimmermann, 1958: 347, as the type species of the genus *Hednota* MEYR. *Hednota* MEYR. was sunk under *Talis* GUEN., by HAMPSON, in 1896: 967. However, the species of the genus *Hednota* MEYR. show closer relationship to the members of

the genus *Chilo* GERM. & ZCK., than to those of the genus *Talis* GUEN., as the preliminary study of the present author has shown. Thus, the genus *Hednota* MEYR. should be validated, but it is in need of thorough revision as, most probably, it comprises a heterogenous cluster of species belonging to several distinct genera.

***Hemiplatytes* BARNES & BENJAMIN**

Hemiplatytes BARNES & BENJAMIN, 1924, Contr. n. H. Lep. N. Amer., 5: 192.

This genus was erected for two species, namely *Diatraea epia* DYAR and *Diatraea prosenes* DYAR. Type species *Diatraea epia* DYAR, 1912: 165, from California, by original designation.

***Hemiptocha* DOGNIN**

Hemiptocha DOGNIN, 1905, Ann. Soc. ent. Belg., 49: 61.

Established for a single species, *Hemiptocha agrapheella* DOGNIN, 1905: 61, Tucuman, South America.

Hombergia J. JOAN., see *Cephis* RAG.

***Hoploscopa* MEYR.**

Hoploscopa MEYRICK, 1886, Trans. ent. Soc. Lond., 1886: 267.

Erected for *Hoploscopa astrapias* MEYR. 1886: 268, from Fiji, in *Scopariidae*. In 1932: 344, MEYRICK transferred this genus to *Crambidae*.

***Hypiesta* HMPS.**

Hypiesta HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 3: 538.

This genus was erected for two species, namely *Hypiesta argyrogramma* HMPS. and *H. flavirupalis* HMPS. Type species, *Hypiesta argyrogramma* HMPS., 1919: 538, by original designation.

***Hypotomorpha* REBEL**

Hypotomorpha REBEL, 1892, Ann. Hofmus. Wien, 7: 252.

Established for a single species, namely, *Hypotomorpha lancerotella* REBEL, 1892: 253, from the Canary Islands.

Iesta DYAR, see *Diatraea* GUILD.

Japonichilo OKANO, 1962

Japonichilo OKANO, 1962, Ann. Rep. Gak. Fac. Iwate Univ., **20**: 121.

Established for a single species, *Japonichilo bleszynskii* OKANO from Japan and China (OKANO, 1962b: 121).

Japonicrambus OKANO, 1962

Japonicrambus OKANO, 1962, Ann. Rep. Gak. Fac. Iwate Univ., **20**: 104.

Established for two species, namely, *Crambus bilineatus* OKANO, 1958: 78 and *Japonicrambus ishizukai* OKANO, 1962b: 104, both from Japan. Type species: *Crambus bilineatus* OKANO, designated by OKANO in the original description of this genus.

Jartheza WALK., see *Ancylolomia* HBN.

Juania AUR.

Juania AURIVILLIUS, 1922, Nat. Hist. Juan Fern. (3) **2**: 264.

Erected for a single species, namely, *Juania annulata* AUR., 1922: 265, from Masatierra, Juan Fernandez.

Kupea PHILPOTT

Kupea PHILPOTT, 1930, Rec. Canterbury Mus., **3**: 247.

Erected for a single species, *Kupea electilis* PHILPOTT, 1930: 247, from New Zealand.

Leucargyra HMPS.

Leucargyra HAMPSON, 1896, Proc. zool. Soc. Lond., **1895**: 960.

Established for one species, namely, *Leucargyra puralis* HMPS., 1896: 960, from Therespolis, Brazil.

Loxocrambus FORBES

Loxocrambus FORBES, 1920, J. N. York ent. Soc., **28**: 225.

Established for two species, namely, *Loxocrambus canellus* FORBES and *L. mohaviellus* FORBES. Type species, *Loxocrambus canellus* FORBES, 1920: 226, from Biloxi, Missouri, by original designation.

***Loxophantis* MEYR.**

Loxophantis MEYRICK, 1935, Exot. Micr., 4: 570.

Erected for a single species, namely *Loxophantis triplecta* MEYR., 1935: 570.

Macrochilo HAMPSON, 1896: 950, erected for *Chilo ambiguellus* SNELL., 1890: 642, is referable to *Schoenobiidae* (BŁESZYŃSKI 1962a: 3). The present author gave a new name *Promacrochilo* BŁESZ. for *Macrochilo* Hmps., nec *Macrochilo* HÜBNER, 1825.

Margaretania AMSEL, 1961, Ark. Zool. (2) 13: 329, erected for a single species, namely *M. superba* AMSEL, 1961: 329 is referable to *Pyraustidae*, *Cybalomiinae*.

***Mesocrambus* BŁESZ.**

Mesocrambus BŁESZYŃSKI, 1957, Acta zool. cracov., 1: 250.

Erected for a single species, namely *Crambus candiellus* H.-S., 1848: 56, from Europe.

***Mesolia* RAG.**

Mesolia RAGONOT, 1888, Ann. Soc. ent. Fr., (6) 8: 282
Eugrotea FERNALD, 1896, Cramb. N. Amer.: 16.

The genus *Mesolia* RAG. was erected for a single species, *Mesolia pandavella* RAG., 1888: 282, from India and Ceylon. The genus *Eugrotea* FERN. was established for *Eugrotea dentella* FERN., 1896: 16, from Florida; it was sunk under *Mesolia olivella* GROTE by FORBES in 1920: 216.

***Mestolobes* BUTL.**

Mestolobes BUTLER, 1882, Trans. ent. Soc. Lond., 1882: 37.

Erected for four species, *Mestolobes aeone* BUTL., *M. semiochrea* BUTL., *M. simaethina* BUTL. and *Metasia abnormis* BUTL., Hawaii. Type species *Metasia abnormis* BUTL., 1882: 35 (ZIMMERMANN, 1958: 301). *Promylaea* MEYR., 1899: 227, erected for a single species, *Promylaea pyropa* MEYR., 1899, is considered by ZIMMERMANN, 1958: 327, as a subgenus of *Mestolobes* MEYR.

***Metacrambus* BŁESZ.**

Metacrambus BŁESZYŃSKI, 1957, Acta zool. cracov., 1: 253.

Erected for a single species, namely, *Crambus carectellus* ZELL., 1847: 751, described from Sicily.

***Metaeuchromius* BŁESZ.**

Metaeuchromius BŁESZYŃSKI, 1960 Acta zool. cracov., 5: 217.

Erected for a single species, namely, *Eromene yuennanensis* CAR. [in] CAR. & MEYR., 1937: 151, from China, Prov. Yuennan.

Meyrickella BERG, 1898, Commun. Mus. Buenos Aires, 1 (2): 42.

This is a new name for the genus *Prionophora* MEYR., 1879: 179, which is praecoccupied by *Prionophora* DEJEAS, 1833, Coleoptera. The genus was erected for *Crambus rupstellus* WALK., 1863:173 (referable to *Noctuidae*).

***Microchilo* OKANO**

Microchilo OKANO, 1962, Ann. Rep. Gak. Fac. Iwate Univ. 20: 129.

Established for one species, *Microchilo inouei* OKANO, 1962b: 129, from Japan and China.

***Microtalis* TURN.**

Microtalis TURNER, 1911, Ann. Queensland Mus., 10: 116.

This genus was established for two species, namely, *Microtalis epimetalla* TURN. and *M. acrocapna* TURN. Type species, *Microtalis epimetalla* TURN., 1911: 116, from Port Darwin, North Australia, by original designation.

Mimopolyocha MATSUMURA, 1925, J. Coll. agric. Hokkaido imp. Univ., 15: 184. Erected for a single species, namely, *Platytes obscurella* MATS., 1911: 57, from Sakhalin. Referable to *Phycitidae*.

***Miraxis* BŁESZ.**

Miraxis BŁESZYŃSKI, 1962, Coridon, (B) 3: 6.

Established for one species, namely, *Miraxis klotzi* BŁESZ., Coridon, 1962, (B) 3: 6, from Peru.

Mithotemma BUTLER, 1883, Trans. ent. Soc. Lond. 1883: 62. Erected in *Crambidae*. HAMPSON, in 1896: 974, transferred this genus to *Noctuidae*.

Mixophyla MEYRICK, 1887, Trans. ent. Soc. Lond. 1887: 268. Erected in *Crambidae* for *Crambus ermineus* MOORE. In 1896: 269 transferred by HAMPSON to *Hydrocampinae*.

Miyakea MARUMO

Miyakea MARUMO, 1933, Classification of the subfamily *Crambinae* in Japan, Summary: 2.

Erected for a single species, namely, *Eromene expansa* BUTL., 1881: 590, from Japan.

Myzea WALK., see *Calamotropha* ZELL.

Monotonia AMSEL, 1955, Bull. Inst. Sci. nat. Belg., 31 (83): 44. Described for *?Adelosomia straminella* ZERNY, 1914: 316, from Near East. Referable to *Phycitidae*.

Myelobia H.—S.

Myelobia HERRICH-SCHAFFER, 1858, Samml. aussereur. Schmett.: 79

Doratoperas HAMPSON, 1896, Proc. zool. Soc. Lond. 1895: 961

Chilopsis HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 4: 56

Xanthopherne DYAR & HEINRICH, 1927, Proc. U. S. nat. Mus., 71 (19): 29.

Myelobia H.-S. was described for *M. pustulata* H.-S., *M. paleacea* H.-S., *M. murina* H.-S. and *M. decolorata* H.-S., all from South America. This genus has hitherto been placed in *Galleriidae* (BŁESZYŃSKI, 1962b: 1). *Doratoperas* HMPS. was erected for a single species, namely, *Crambus? atrosparsellus* WALK., 1863: 163, from Colombia and Brazil. *Chilopsis* HMPS. was established also for a single species, namely *Ch. squamata* HMPS. 1919: 56, from Peru. *Xanthopherne* DYAR & HEINRICH was described for two species, namely, *Doratoperas biumbrata* SCHAUS and *D. fulvescens* HMPS. The latter is referable to the genus *Haimbachia* DYAR and the former was designated originally as the type species of *Xanthopherne* DYAR & HEINR. Type species, *M. paleacea* H.-S. (BŁESZYŃSKI & COLLINS, 1962: 326).

Myzea WALK., see *Calamotropha* ZELL.

Myriostephes MEYR.

Myriostephes MEYRICK, 1884, Trans. ent. Soc. Lond., 1884: 277.

Erected for a single species, *Myriostephes matura* MEYR., 1884: 328, in *Pyraustidae*. According to TURNER, referable to *Crambidae*.

Naurace WALK., see *Prionapteryx* STEPH.

Nagahama MARUMO, see *Platytes* GUEN.

Neargyria HMPS.

Neargyria HAMPSON, 1896, Proc. zool. Soc. Lond., 1895: 923.

Erected for a single species, namely *Argyria argyraspis* MEYR., 1879, Proc. Linn. Soc. N. S. Wales, 4: 216, from Australia.

Neerupa HMPS.

Neerupa HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 3: 278.

Erected for two species, namely *Neerupa benepunctalis* HMPS., and *N. argyrosticta* HMPS., from Colombia. Type species *Neerupa benepunctalis* HMPS., 1919: 278, by original designation.

Neocrambus BŁESZ.

Neocrambus BŁESZYŃSKI, 1957, Acta zool. cracov., 1: 457.

Erected for a single species, namely *Crambus wolfschlägeri* SCHAW., 1937, Z. Öst. ent. Ver., 22: 55, from Macedonia, South Europe.

Nephalia TURN.

Nephalia TURNER, 1911, Ann. Queensland Mus., 10: 113.

Erected for a single species, namely *Nephalia crypsimetalla* TURN., 1911: 114, from Australia.

Notocrambus TURN.

Notocrambus TURNER, 1922, Proc. Roy. Soc. Vict., 35: 45.

Erected for a single species, namely, *Notocrambus holomelas* TURN., 1922: 46, from Tasmania.

Novocrambus AMSEL

Novocrambus AMSEL, 1956, Bol. Ent. Venezolana, 10: 30.

Erected for one species, namely *Crambus pygmaeus* ZELL., 1881: 173, from Colombia. This species has previously been placed in the genus *Ptochostola* MEYR. Actual name of *C. pygmaeus* ZELL. (nec *C. pygmaeus* STEPHENS, 1834, is *Novocrambus propygmaeus* BŁESZ. & COLLINS, 1962: 328.

Occidentalia DYAR & HEINR.

Occidentalia DYAR & HEINRICH, 1927, Proc. U. S. nat., Mus. 71 (19): 41.

This genus was established for a single species, namely *Crambus comptulatalis* HULST, 1886: 167.

Ommatopteryx KIRBY, see *Euchromius* GUEN.

Oressaula TURN., see *Tauroscopa* MEYR.

***Orocrambus* PURDIE**

Orocrambus PURDIE, 1884, N. Zeal. J. Sci., 2: 167.

Orocrambus MEYRICK, 1885, Trans., N. Zeal. Inst., 17: 133.

Erected for a single species, *Orocrambus melampetrus* PURDIE, 1883: 168, from New Zealand. This genus was discovered and described by MEYRICK, but PURDIE's description is earlier and valid.

***Orthomecyna* BUTL.**

Orthomecyna BUTLER, 1883, Ent. mon. Mag., 19: 178.

Erected for *Mecyna exigua* BUTL., 1879: 271, and *Orthomecyna albicaudata* BUTL., 1883: 178. Type species *Mecyna exigua* BUTL., from Hawaii, by original designation.

***Pachymorphus* MÖSCHLER**

Pachymorphus MÖSCHLER, 1890, Abh. Senck. Ges., 16: 324.

This genus was erected for a single species, namely *Pachymorphus subductellus* MÖSCHL., 1890: 324, from Porto Rico.

***Pagmania* AMSEL**

Pagmania AMSEL, 1961, Ark. Zool., (2) 13: 332.

Erected for a single species, namely *Pagmania bilinealis* AMSEL, 1961: 332, from Iran.

Palparia HAW., see *Crambus* F.

***Parancyla* HMPS.**

Parancyla HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 4: 67.

This genus was described for a single species, namely *Parancyla argyrothysana* HMPS., 1919: 67, from Central Africa.

***Paratraea* HMPS.**

Paratraea HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 3: 542.

Established for two species, namely *Paratraea plumbipicta* HMPS. and *P. griseifasciata* HMPS., from Transvaal. Type species *Parathraea plumbipicta* HMPS., 1919: 542, by original designation.

Parerupa Hmps.

Parerupa HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 3: 539
Coenotalis HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 4: 152.

Parerupa Hmps. was described for one species, namely *Parerupa diagonalis* Hmps., 1919: 539, from East Africa. This species was sunk under *Diatraea africana* AUR. 1910: 54, from Kilimandjaro, Africa, by MARTIN, in 1958: 190. The genus *Coenotalis* Hmps. was described for a single species, namely *Coenotalis distictalis* Hmps., 1919: 152, from Nigeria. The present writer, in 1962a: 2, sank *Coenotalis* Hmps. under *Parerupa* Hmps.

Pareromene Osth.

Pareromene OSTHEIDER, 1941, Mitt. Münch. ent. Ges. 31: 366.

Erected for a single species, namely *Pareromene rebeli* Osth., 1941: 366, from Crete.

Pediasia Hbn.

Crambus FABRICIUS, 1798, Ent. Syst. Suppl.: 464 (in part)
Palparia HAWORTH, 1811, Lep. Brit., (3): 481 (in part)
Chilo ZINCKEN, 1817, Mag. Ent., 2: 34 (in part)
Pediasia HÜBNER, [1825], Verz. bek. Schmett.,: 365
Carvanca WALKER, 1856, List. Spec. lep. Ins. B. M., 9: 119.

The genus *Pediasia* Hbn. was erected for six species, of which four are actual members of this genus and two others belong to the genus *Agriphila* Hbn. The type species is *Tinea fascelinella* Hbn., [1813], fig. 368, Europe; it was designated by the writer in 1953: 102. *Carvanca* WALK. was described for a single species *C. trisecta* WALK., 1856: 119, from North America, was synonymized with *Crambus* F. by HAMPSON 1896: 925. However, *C. trisecta* WALK. is congeneric with *P. fascelinella* Hbn., and falls to the synonyms of *Pediasia* Hbn.

Phanerobela Turn.

Phanerobela TURNER, 1932, Trans. roy. Soc. South Austr. 56: 189.

Erected for a single species, namely *Phanerobela niphospila* TURN., 1932: 189, from Dunk Island, Australia.

***Phanomorpha* TURN.**

Phanomorpha TURNER, 1937, Proc. roy. Soc. Queensland, **48**: 65.

Erected for a single species, namely *Phanomorpha leucoxantha* TURN., 1937: 65, from Queensland, Australia.

Pindicitora WALK., see *Surattha* WALK.

***Platytes* GUEN.**

Palparia HAWORTH, 1811, Lep. Brit., **3**: 481 (in part)
Chilo ZINCKEN, 1817, Mag. Ent., **2**: 34 (in part)
Argyroteuchia HÜBNER, [1825], Verz. bek. Schmett.: 366 (in part)
Thisanotia HÜBNER, [1825], Verz. bek. Schmett.: 366 (in part)
Araxes STEPHENS, 1834, Ill. Brit. Ent. Haust., **4**: 315 (in part)
Platytes GUENÉE, 1845, Ann. Soc. ent. Fr., (2) **3**: 324
Nagahama MARUMO, 1933, Classif. Sumbfam. Cramb. Jap., Summary: 2.

This genus was erected for one species, namely *Tinea cerussella* D. & SCHIFF. HAMPSON in 1896: 943 gives in the synonymy of *Platytes* GUEN. the following genera: *Argyria* HBN., *Catharylla* ZELL., *Urola* WALK. and *Ptychopseustis* MEYR. However, all these genera except *Urola* WALK. are valid. Thus, the genus *Platytes* GUEN. sensu HAMPSON, was a heterogeneous cluster of species. HAMPSON, in 1919, considered the name *Platytes* GUEN. to be a nomen nudum, but such an opinion is incorrect as the GUENÉE genus is well indicated by two species given under it. *Platytes* GUEN. seems to comprise still a heterogeneous complex of species, many of which need to be placed in genera of their own. As far I know, the true members of *Platytes* GUEN. occur only in the Palaearctic Region. The genus *Nagahama* MAR. erected for a single species, *Crambus ornatellus* LEECH, from Japan, was synonymized with *Platytes* GUEN. (BŁESZYŃSKI, 1962d: 25).

Platytesia STRAND, see *Surattha* WALK.

***Polyterpnes* TURN.**

Polyterpnes TURNER, 1932, Trans. roy. Soc. South Austr., **56**: 188.

Established for a single species, namely *Polyterpnes polyrrhoda* TURN. 1932: 189, from North-western Australia.

***Prionapteryx* STEPH.**

Prionapteryx STEPHENS, 1834, Ill. Brit. Ent. Haust., **4**: 317
Naurace WALKER, 1863, List. Spec., **27**: 188.

Erected for a single species, namely *Prionapteryx nebulifera* STEPH., 1834: 317, Florida. The genus *Naurace* WALK. was erected for a single species, *Naurace eugraphis* WALK. 1863: 188, from San Domingo; it was sunk under *Prionapteryx* STEPH., by HAMPSON, in 1896: 963.

Prionophora MEYR., see *Meyrickella* BERG.

Prionotalis HMPS.

Prionotalis HAMPSON, 1919, Ann. Mag. nat. Hist., (9) 4: 152.

Erected for a single species, namely *Prionotalis peracutella* HMPS., 1919: 153, from Gold Coast and British Central Africa.

Proceras BOJER

Proceras BOJER, 1856, Rapport of the Committee on the „Cane Borer.“ Mauritius, 4 plates
Borer GUENÉE in MAILLARD, 1862, Notes sur l'Île de la Réunion par L. Maillard. Lépidoptères. G 70.

The genus *Proceras* BOJER included a single species, namely, *Proceras sacchariphagus* BOJER, 1856, from Mauritius. The genus *Borer* GUEN., was erected for one species, *Borer saccharellus* GUEN., 1862: G70, from Mauritius. The revival of the genus *Proceras* BOJER was proposed by TAMS, 1942: 67. The same author sank the GUENÉE genus under *Proceras* BOJER.

Productalius MARION

Productalius MARION, 1954, Mém. Inst. sci. Madagascar (E) 5: 49.

Erected for a single species, namely *Productalius tritaeniellus* MARION, 1954: 49, from Madagascar.

Promylaea MEYR., see *Mestolobes* BUTL.

Propexus GROTE, see *Thaumatopsis* MORR.

Prosmixis ZELL., see *Talis* GUEN.

Protaphomia MEYR.

Protaphomia MEYRICK, 1936, Arb. morph. taxon. Ent., 3: 97.

According to a kind information of Mr. P. E. S. WHALLEY of the British Museum (N. H.), London, the genus *Protaphomia* MEYR., described in *Galleriidae*, is referable to *Crambidae*, and probably congeneric with *Myelobia* H.-S. This genus was described for one species, viz. *Protaphomia haplodoxa* MEYR., 1936: 97, from Brazil.

Protyparcha MEYR.

Protyparcha MEYRICK, 1909, Subantarct. Isl. New Zealand, 1: 71.

Erected for a single species, namely *Protyparcha scaphodes* MEYR., 1909: 71, from Auckland Island.

Pseudargyria OKANO

Pseudargyria OKANO, 1962, Trans. lep. Soc. Japan, 12: 51.

Erected for a single species, *Argyria interruptella* WALK., 1866: 1763, from Japan and China.

Pseudobissetia BŁESZ.

Pseudobissetia BŁESZYŃSKI, 1959, Beitr. natur. Forsch. Südwestdeutschl., 18 (1): 115.

Erected for a single species, namely *Chilo terrestrellus* CHRIST., 1885: 151.

Pseudocatharylla BŁESZ.

Urola WALKER, 1863, List. Spec., 27: 181 (in part)

Catharylla ZELLER, 1863, Chil. Cramb. Gen. Spec.: 50 (in part)

Pseudocatharylla BŁESZYŃSKI, 1961, Zeit. Wien. ent. Ges., 46: 33.

This genus was erected for four species, namely *Crambus flavoflabellus* CAR., *Crambus inclaralis* WALK., *Crambus aurifimbriellus* HMPS. and *Crambus infixellus* WALK. Generic type is *Crambus flavoflabellus* CAR. by the original designation. However, *Crambus flavoflabellus* CAR., 1925: 298, described from Shanghai, is a synonym of *Crambus inclaralis* WALK. as a recent study of the types of the two species has shown. The species cited by the present writer (1961c: 35) as *Pseudocatharylla inclaralis* (WALK.) was in fact *P. simplex* (ZELL.), which is also a typical member of the genus under consideration.

Pseudoctenella STRAND, see *Ancylolomia* HBN.

Pseudometachilo BŁESZ.

Pseudometachilo BŁESZYŃSKI, 1962, Pol. Pis. ent., 32: 1.

Erected for a single species, namely *Crambus diatraeellus* (HMPS.), 1896: 931 (= *Crambus distictellus*, HMPS., 1896: 938), from Brazil and Argentina.

Pseudoschoenobius FERN.

Pseudoschoenobius FERNALD, 1896, Cramb. N. Amer.: 18.

Erected for one species, viz. *Schoenobius opalescalis* HULST, 1886: 167, from Arizona.

Ptochostola MEYR.

Ptochostola MEYRICK, 1882, Proc. Linn. Soc. N. S. Wales, 7: 154.

Erected for a single species, namely *Crambus dimidiellus* MEYR., 1879: 190, from Sydney (synonym of *Crambus microphaeellus* WALK., 1866: 1758).

Ptychopseustis MEYRICK, 1889, Trans. ent. Soc. Lond., 1889: 521. Described in *Crambidae* for *Diptychophora amoenella* SNELL, 1880: 247. Referable to *Pyraustidae* (BŁESZYŃSKI, 1962a: 3).

Raphiptera HMPS.

Raphiptera HAMPSON, 1896, Proc. zool. Soc. Lond. 1895: 961.

Erected for a single species, namely *Crambus minimellus* ROB., 1871: 315, from Pennsylvania, North America.

Scenoploca MEYR.

Scenoploca MEYRICK, 1882, N. Zeal. J. Sci. 1: 186.

Established for one species, viz. *Scenoploca petraula* MEYR., 1882: 186, from New Zealand.

Sedenia GUENÉE, 1854, Delt. & Pyr.: 249. Described in *Crambidae* but referable to *Pyraustidae* (BŁESZYŃSKI, 1962a: 3).

Sericocrambus WALL.

Sericocrambus WALLENGREN, 1868, Eugenies Reise, Lep.: 383.

Erected for a single species, *Sericocrambus stylatus* WALL., 1868: 383, from Argentina.

Silveria DYAR, see *Chilo* ZCK.

Soroscotia ROSENSTOCK, 1885, Ann. Mag. nat. Hist., (5) 16: 435. Described for *?Eromene vetustella* WALK., from Australia. Referable to the family *Nolidae*.

***Sphaerodeltis* MEYR.**

Sphaerodeltis MEYRICK, 1933, Exot. Mier., 4: 380.

Erected for a single species, namely *Sphaerodeltis psammoleuca* MEYR., 1933: 380, from Congo.

Stenochilo HMPS., see *Cephis* RAG.

***Styxon* BŁESZ.**

Styxon BŁESZYŃSKI, 1962, Pol. Pis. ent., 32: 6.

This genus was erected for a single species, *Stenochilo ciniferalis* CAR., 1925, Mem. Sect. științ. Acad. rom., (3) 3: 299, described from Kwangtung, China.

***Surattha* WALK.**

Surattha WALKER, 1863, List Spec., 27: 75

Pindicitora WALKER, 1863, List Spec., 27: 134 (in part)

Calarina WALKER, 1866, List Spec., 35: 1769

Platytesia STRAND, 1919, Ent. Ztg. Stettin, 79: 255.

The genus *Surattha* WALK. was established for a single species, namely *Surattha invectalis* WALK., 1863: 76, from India. The genus *Pindicitora* WALK. was erected for four species, of which, *Pindicitora thysbesalis* WALK., 1863: 135, is a synonym of the preceding species. (HAMPSON, 1896: 965). The generic type of the genus *Pindicitora* WALK. is *P.zeuxoalis* WALK., designated by WARREN, 1892: 172, but this species is not referable to *Crambidae*. The genus *Calarina* WALK. was described for one species, *Calarina albirenella* WALK., 1866: 1770, from India and Java. *Platytesia* STRAND was erected as a subgenus of *Platytes* GUEN., for one species, namely *Platytesia alikangiella* STRAND, from Formosa. SHIBUYA, in 1928: 14, sank this species under *Surattha albistigmata* WIL. & SOUTH 1918: 217, also from Formosa.

Syncrotaula MEYRICK, 1933, Exot. Mier., 4: 378 (= *Eudorina* SNELLEN. 1895, Tijd. Ent., 38: 116, nec *Eudorina* EHRENBURG, 1832, in *Protozoa*). Hitherto placed in *Crambidae*, but referable to *Pyraustidae* (BŁESZYŃSKI, 1962a: 3).

***Talis* GUEN.**

Talis GUENÉE, 1845, Ann. Soc. ent. Fr., (2) 3: 324

Prosmixis ZELLER, 1846, Linn. Ent., 1: 270

Araxates RAGONOT, 1888, Ann. Soc. ent. Fr., (6) 8: 281.

The genus *Talis* GUEN. was erected for a single species, namely *Tinea querella* DEN. & SCHIFF., 1775: 134, Europe. HAMPSON, in 1919: 147, sank *Talis* GUEN. under *Prosmixis* ZELL., considering the former name to be a nomen nu-

dum. However, such an opinion is incorrect as the genus *Talis* GUEN. was well indicated by a species placed under it. The genus *Araxates* RAG. was erected for a single species, namely, *Crambus pulcherrimus* STGR., 1870: 192, Europe. HAMPSON, in 1896: 967, sank the RAGONOT name under *Talis* GUEN. The genus *Hednota* MEYR. sunk by HAMPSON, 1896: 967, under *Talis* GUEN. should be validated as close to *Chilo* ZCK. and quite different from *Talis* GUEN. (discussion see above).

***Tamsica* ZIMM.**

Tamsica ZIMMERMANN, 1958, Ins. Haw., 8: 346.

This genus was erected for six Hawaiian species, hitherto placed in the genus *Talis* GUEN., or *Hednota* MEYR. Type species *Gesneria floricolens* BUTL., 1883: 180, by original designation.

Taseopteryx BUTLER, 1883, Trans. ent. Soc. Lond. 1883: 63.

Erected for a single species, namely *Taseopteryx sericea* BUTL., 1883: 64, from Chili. Referable to *Torticidae*.

***Tauroscopa* MEYR.**

Tauroscopa MEYRICK, 1888, Trans. N. Zeal. Inst., 20: 69.

Erected for a single species, namely *Tauroscopa gorgopis* MEYR., 1888: 69, from New Zealand.

***Tawhitia* PHILP.**

Tawhitia PHILPOTT, 1931, Trans. N. Zeal. Inst., 62: 26.

Erected for a single species *Tauroscopa glaucophanes* MEYR., 1907, Trans. N. Zeal. Inst., 39: 110.

***Tehama* HULST**

Tehama HULST, 1888, Ent. Amer., 4: 115.

Described for one species, namely *Spermatophthora bonifatella* HULST, 1887: 135, from North America.

***Thalamarchis* MEYR.**

Thalamarchis MEYRICK, 1897, Trans. ent. Soc. Lond., **1897**: 80.

Erected for a single species, namely *Thalamarchis chalchorma* MEYR., 1897: 81, from Sangir, Australian Region.

***Thaumatopsis* MORR.**

Thaumatopsis MORRISON, 1874, Proc. Boston Soc. nat. Hist., **17**: 161.

Propexus GROTE, 1880, Can. Ent., **13**: 19.

The genus *Thaumatopsis* MORR. was erected for one species, namely, *Thaumatopsis longipalpus* MORR., 1874: 165, from North America. However, this species is a synonym of *Crambus pexellus* ZELL., 1863: 48 (FERNALD, 1896: 66). The name *Propexus* was used by GROTE who described (1880: 19) a species under a name *Crambus Propexus edonis* GROTE. Both, *Crambus pexellus* ZELL. and *Crambus Propexus edonis* GROTE are congeneric (FERNALD, 1896: 65).

***Thisanotia* HBN.**

Crambus FABRICIUS, 1798, Ent. Syst. Suppl.: 464 (in part)

Palparia HAWORTH, 1811, Lep. Brit., (3): 481 (in part)

Chilo ZINCKEN, 1817, Mag. Ent., **2**: 34 (in part)

Thisanotia HÜBNER, [1825], Verz. bek. Schmett.: 366.

The genus *Thisanotia* HBN. was erected for three species, namely *Tinea chrysonuchella* SCOP., *Tinea rorella* L. and *Tinea auriferella* DEN. & SCHIFF. The species *Tinea chrysonuchella* SCOP., 1763: 246, Europe, was designated by the present writer in 1957: 182, as the type species of the genus in question. All three species, which were placed under *Thisanotia* HBN., are also members of the HAWORTH *Palparia* HAW. The name *Thisanotia* HBN. was adopted by HEINEMANN, 1863: 116 for two species, *T. cerussella* (DEN. & SCHIFF.) and *T. alpinella* (HBN.). The two are actual members of the genus *Platytes* GUEN. HEINEMANN used an incorrect spelling of the HÜBNER name, writing *Thinasotia* instead of *Thisanotia*. This error has subsequently been repeated by MEYRICK, who adopted the HÜBNER name for several Australian species, which are actually members of the genus *Hednota* MEYR.

Tollia AMSEL, see *Ancylolomia* HBN.

***Tulla* ZIMM.**

Tulla ZIMMERMANN, 1958, Ins. Haw., **8**: 336.

Erected for a single species, namely, *Prionapteryx exonoma* MEYR., 1899: 198, Oahu, Hawaii.

***Ubida* WALK.**

Ubida WALKER, 1863, List Spec., 27: 185

Crunophila MEYRICK, 1882, Proc. Linn. Soc. N. S. Wales, 7: 152.

The genus *Ubida* WALK. was erected for a single species, namely *Ubida receptalis* WALK., 1863: 186, from Moreton Bay, Australia. The genus *Crunophila* MEYR. was described for *Crambus ramosstriellus* WALK., 1863: 172, from Australia. BUTLER, in 1884: 75, pointed out that *Crunophila* MEYR. is a synonym of *Ubida* WALK. as *Crambus ramosstriellus* WALK. is conspecific with *Ubida receptalis* WALK.

***Uinta* HULST**

Uinta HULST, 1888, Ent. Am., 4: 116.

Erected for a single species, namely *Uinta oreadella* HULST, 1888: 116, from Colorado, North America.

Urola WALK., see *Argyria* HBN.,

Uscodys DYAR, 1908, Proc. ent. Soc. Wash., 10: 101.

Erected for a single species, namely *Anerastia cestalis* HULST, 1886: 164. Referable to *Noctuidae*.

***Vaxi* BŁESZ.**

Vaxi BŁESZYŃSKI, 1962, Pol. Pis. ent., 32: 11.

Described for a single species, namely *Conocrambus obliqua* HMPS., 1919: 444, from Brazil. Close to the genus *Argyria* HBN. The male of *Vaxi obliqua* (HMPS.) is unknown.

***Velasquez* BŁESZ.**

Velasquez BŁESZYŃSKI, 1962, Pol. Pis. ent., 32: 14.

Established for *Crambus pentadactylus* ZELL., 1863: 38 (= *Aquita clavifarella* WALK., = *Aphomia strigosa* BUTL.) from New Zealand and Tasmania.

***Veronese* BŁESZ.**

Veronese BŁESZYŃSKI, 1962, Pol. Pis. ent., 32: 27.

Established for one species, namely, *Crambus distinctellus* LEECH, 1889: 107, from East Asia.

Xanthocrambus BŁESZ.

Xanthocrambus BŁESZYŃSKI, 1955, Zeit. Wien. ent. Ges. **40**: 266.

Described for three species, namely, *Crambus delicatellus* ZELL., *Chilo saxo-nellus* ZCK. and *Crambus occidentellus* CAR. Type species *Crambus delicatellus* ZELL., 1863: 41, from South Europe, by original designation.

Xanthopherne DYAR & HEINRICH, see *Myelobia* H.-S.

Xubida SCHAUS

Xubida SCHAUS, 1922, Proc. U. S. nat. Mus., **24**: 140.

Erected for twelve species; of these, five species were described as new and the remaining ones were transferred from the genera *Platytes* GUEN. and *Ubida* WALK. The type species, *Xubida dentilineella* SCHAUS, 1922: 141, from Guatemala, by original designation.

Zacatecas BŁESZ.

Zacatecas BŁESZYŃSKI, 1962, Pol. Pis. ent., **32**: 6.

Erected for a single species, namely *Crambus ankasokellus* VIETTE, 1960, Rev. fr. d'Ent., 27 (2): 200, from Madagascar.

Zeadiatraea Box

Zeadiatraea Box, 1955, Proc. R. ent. Soc. Lond., (B) **24**: 197.

Erected for four species, namely, *Leucania lineolata* WALK., *Diatraea grandiosella* DYAR, *D. schausella* DYAR & HEINR. and *D. muellerella* DYAR & HEINR. Type species *Leucania lineolata* WALK., 1856: 100, from South America, by original designation.

Zolea WALK., see *Erupa* WALK.

Zovax BŁESZ.

Zovax BŁESZYŃSKI, 1962, Acta zool. cracov., **7**: 130.

Established for one species, namely *Prionopteryx* [sic] *whiteheadi* WOLASTON, 1879, Ann. Mag. nat. Hist., (5) **3**: 340, from Santa Helena.

REFERENCES

- AMSEL H. G. 1935. Neue palästinische Lepidopteren. Mitt. zool. Mus., Berlin, **20**: 271—319
18 pls.
- AMSEL H. G. 1949. On the *Microlepidoptera* collected by E. P. WILTSHERE in Irak and Iran,
in the years 1935- to 1938. Bull. Soc. Fouad I Ent., Cairo, **33**: 271—351, 12 pls.
- AMSEL H. G. 1955. Über mediterrane Microlepidopteren und einige transkaspische Arten.
Bull. Inst. Sci. nat. Belg., Bruxelles, **31**, (83) 64 pp., 6 pls.
- AMSEL H. G. 1956-1957. *Microlepidoptera* Venezolana I, II. Bol. Ent. Venez. Maracay, **10**:
1—336 (1956); II, 110 pls. (1957).
- AMSEL H. G. 1961. Die Mierolepidopteren der BRANDT'SCHEN Iran-Ausbeute. 5. Teil. Ark.
Zool., Stockholm, (2) **13** 323—445, 9 pls., 136 figs.
- AURIVILLIUS C. 1910. *Lepidoptera*. In: SjÖSTEDTS Kilimandjaro-Meru Expedition Stockholm.
9: 1—56, 2 pls.
- AURIVILLIUS C., PROUT L. B., MEYRICK E. 1922. Lepidopteren von Juan Fernandez und der
Oster-Insel. Nat. Hist. Juan Fernandez, Uppsala, **3**: 255—270, pls. x—xi.
- BARNES W., BENJAMIN F. H. 1924. Nomenclature. New Species. Contr. nat. Hist. Lep. N. Amer.,
Illinois, **5**: 99—199.
- BERG C. 1898. Substitución de nombres genericos. Com. Mus. Buenos Aires, Buenos Aires,
1: 40—43.
- BŁESZYŃSKI S. 1954. Studies on the *Crambidae* (*Lepidoptera*). vi. Three new Palearctic species
of the genus *Pediasia* HBN. Ann. Mus. zool. Pol., Warszawa, **15**: 101—108, pls. xv—xvi.
- BŁESZYŃSKI S. 1955. Materialien zur Kenntnis der *Crambidae*. Teil x. Über die systematische
Stellung der *Crambus delicatellus* ZELL. — Gruppe. Zeit. Wien. ent. Ges., Wien, **40**: 266—269.
- BŁESZYŃSKI S. 1956. Materialien zur Kenntnis der *Crambidae*. Teil xiii. Über die Gattung
Catoptria HBN. und eine neue Art dieser Gattung. Zeit. Wien. ent. Ges., Wien, **41**: 213—216,
pl. xxi.
- BŁESZYŃSKI S. 1957. Studies on the *Crambidae* (*Lepidoptera*). Part. xiv. Revision of the Euro-
pean species of the Generic Group *Crambus* F. s. 1. Acta zool. crac., Kraków, **1**: 161—462,
incl. pls. xxvii—xcii.
- BŁESZYŃSKI S. 1959a. Studies on the *Crambidae*. Part xix. Notes on some Japanese species
of the generic group *Crambus* F. Tinea, Tokyo, **5**: 273—276, pl. xxxvi.
- BŁESZYŃSKI S. 1959b. Materialien zur Kenntnis der *Crambinae*. 21. Teil. Studien über einzelne
Crambinae aus dem Nahen Osten (*Lepidoptera: Pyralidae*). Beitr. natur. Forsch. Süd-
westdeutschl., Karlsruhe, **18**: 111—116, pl. iii.
- BŁESZYŃSKI S. 1960a. Studies on the *Crambidae*. Part xxviii. A New Genus *Classeya* for Some
Species of the Genera *Argyria* HBN. and *Platytes* GN. and a New Species of this Genus
from Senegal (*Lepidoptera*). Čas. Česk. Spol. Ent., Praha, **57**: 267—274, 1 pl.
- BŁESZYŃSKI S. 1960b. Studies on the *Crambidae*. (*Lepidoptera*). Part xxix. On Some Species
of the Genus *Euchromius* GN. Acta zool. crac., Kraków, **5**: 203—247, incl. pls. xxi—xxxiv.
- BŁESZYŃSKI S. 1961a. Revision of the World Species of the Family *Crambidae*. Part I. Genus
Calamotropha ZELL. Acta zool. crac., Kraków, **6**: 137—272, 1 fig., pls. xx—lxxi.
- BŁESZYŃSKI S. 1961b. Studies on the *Crambidae*. Part xxx. On Several Species of the Generic
Group *Crambus* F. from the Ethiopian Region with the Descriptions of New Species and
Genera. Pol. Pis. ent., Wrocław, **31**: 165—218, 20 pls.
- BŁESZYŃSKI S. 1961c. Materialien zur Kenntnis der *Crambidae*. Teil xxxi. Bemerkungen über
einige Arten mit, Beschreibungen von zwei neuen Gattungen und einer neuen Art. Zeit.
Wien. ent. Ges., Wien, **46**: 33—38.
- BŁESZYŃSKI S. 1961d. Studies on the *Crambidae* (*Lepidoptera*). Part xxxiii. Preliminary
Report on the Genus *Eoreuma* ELY. Coridon, London, (A) **2**, 4 pp.
- BŁESZYŃSKI S. 1962a. Studies on the *Crambidae* (*Lepidoptera*). Part xxxiv. Miscellaneous
notes. Coridon, London. (A) **3**, 3 pp.

- BŁESZYŃSKI S. 1962b. Studies on the *Crambidae (Lepidoptera)*. Part xxxv. Note on the genus *Morpheis* HBN. Coridon, London, (A) 4, 2 pp.
- BŁESZYŃSKI S. 1962c. Studies on the *Crambidae (Lepidoptera)*. Part. xxxvi. On Some Species of the Genus *Chilo* ZINCKEN. Acta zool. cracov., 7: 107—133, pl. xiii.
- BŁESZYŃSKI S. 1962d. Studies on the *Crambidae (Lepidoptera)*. Part xxxvii. Changes in the Nomenclatory of Some *Crambidae* with the Descriptions of New Genera and Species. Pol. Pis. ent., Wrocław, 32: 5-48, 7 pls.
- BŁESZYŃSKI S. 1962e. Studies on the *Crambidae*. Part 38. Descriptions of A New Genus and Three New Species. Coridon, London, (B) 3: 6-7.
- BŁESZYŃSKI S. & COLLINS J. R. 1962. A Short Catalogue of the World Species of the Family *Crambidae (Lepidoptera)*. Acta zool. cracov., 7: 197—389.
- BOJER W., et al. 1856. Report of the Committee on the „Cane Borer.“ Mauritius.
- BOX H. E. 1931. The Crambine genera *Diatraea* and *Xanthopherne* (*Lep. Pyral.*). Bull. ent. Res., London, 22: 1—50, 5 pls.
- BOX H. E. 1953. New *Crambinae* genera allied to *Diatraea* GUILDFING (*Lepidoptera: Pyralidae*). i. Proc. Roy. ent. Soc. Lond., London, (B) 22: 178—180.
- BOX H. E. 1955. New *Crambinae* genera allied to *Diatraea* GUILDFING (*Lepidoptera: Pyralidae*). iii. Proc. Roy. ent. Soc. Lond., London, (B) 24: 197—200.
- BOX H. E., CAPPS H. W. 1955. New Crambine genera allied to *Diatraea* GUILDFING (*Lepidoptera: Pyralidae*). i (Supplementary note) and ii. Proc. Roy. ent. Soc. Lond., London, (B) 24: 174—178 1 pl.
- BUTLER A. G. 1879. On Heterocerous *Lepidoptera* collected in the Hawaiian Islands by the Rev. T. BLACKBURN. Ent. mon. Mag., London, 15: 269—273.
- BUTLER A. G. 1881. Descriptions of new Genera and Species of Heterocerous *Lepidoptera* from Japan. Trans. ent. Soc. Lond., London, 1881: 579—600.
- BUTLER A. G. 1882. On a small collection of *Lepidoptera* from the Hawaiian Islands. Trans. ent. Soc. Lond., London, 1882: 31—45.
- BUTLER A. G. 1883. On a small series of *Lepidoptera* from the Hawaiian Islands. Ent. mon. Mag., London, 19: 176—180.
- CARADJA A. 1925. Ueber Chinas Pyraliden, Tortriciden, Tineiden nebst kurzen Betrachtungen, zu denen das Studium dieser Fauna Veranlassung gibt. (Eine biogeographische Skizze). Mem. Sect̄. științ. Acad. rom., Bucarest, (3) 3: 257—383, 2 pls.
- CHRISTOPH H. 1885. *Lepidoptera* aus dem Achal-Tekke-Gebiete. Mém. Lép. Rom. 2: 119—171, pls. vi—viii, xv.
- DENIS J. N. C. M., SCHIFFERMÜLLER J. 1775. Ankündigung eines systematischen Werkes von den Schmetterlingen der Wienergegend. Wien, 322 pp., 2 pls.
- DOGNIN P. 1905. Hétérocères nouveaux de l’Amerique du Sud. Ann. Soc. ent. Belg., Bruxelles, 49: 61—90.
- DUPONCHEL P. A. J. 1836. In: J. B. GODART, Histoire Naturelle des Lépidoptères ou Papillons de France. Vol. 10. Paris, 402 pp.
- DYAR H. G. 1909. New species of American *Lepidoptera*. Proc. ent. Soc. Wash., Washington, 11: 19—29.
- DYAR H. G. 1910. Descriptions of Some new species and Genera of *Lepidoptera* from Mexico. Proc. U. S. nat. Mus., Washington, 38: 229—273.
- DYAR H. G. 1911. The American species of *Diatraea* GUILDFING. Ent. News, Philadelphia, 22: 199—207.
- DYAR H. G. 1912. *Lepidoptera* collected at Laguna beach, California. Rep. Laguna Mar. Lab., 1: 163—166.
- DYAR H. G. 1914. Descriptions of new species and genera of *Lepidoptera* from Mexico. Proc. U. S. nat. Mus., Washington, 47: 365—409.
- DYAR H. G. 1925. Some new American moths (*Lepidoptera*). Insec. inscit. menstr., Washington, 13: 1—19.

- DYAR H. G., HEINRICH C. 1927. The American moths of the genus *Diatraea* and allies. Proc. U. S. nat. Mus., Washington, **71** (19), 48 pp., 20 pls.
- ELY C. R. 1910. New *Phycitinae* and *Crambinae*. Proc. ent. Soc., Washington, **12**: 202—204.
- FABRICIUS J. C. 1794. Entomologia Systematica emendata et aucta. 3(2). Hafniae, 349 pp.
- FABRICIUS J. C. 1798. Supplementum Entomologiae Systematicae. Hafniae, 572 pp.
- FERNALD C. H. 1888. On the genus *Diatraea*. Ent. Am., Philadelphia, **4**: 119—120.
- FERNALD C. H. 1896. The *Crambidae* of North America. Massachusetts, 81 pp., 6 pls.
- FISCHER EDLEN VON RÖSSLERSTAMM J. E. 1834. Abbildungen zur Beschreibungen und Ergänzung der Schmetterlingskunde Besonders der Microlepidopterologie als Supplement zu TREITSCHKE's und HÜBNER's europäischen Schmetterlingen, mit erläuterndem Text. Leipzig, 308 pp., 100 pls.
- FORBES W. T. M. 1920. Notes on the *Crambidae* (*Lepidoptera*). J. New York ent. Soc., New York, **28**: 214—227.
- GROTE A. R. 1880. *Crambidae*. Can. Ent., London, Ont., **12**: 15—19.
- GROTE A. M. 1882. North American Moths, with a preliminary catalogue of the species of *Hadena* and *Polia*. Bull. U. S. geol. Surv., Washington, **6**: 257—277.
- GUENÉE M. A. 1845. Essai sur une nouvelle classification des Microlépidoptères. Ann. Soc. ent. Fr., Paris, (2) **3**: 297—244.
- GUENÉE M. A. 1854. Historie Naturelle des Insectes. Species Général des Lépidoptères. Par M. M. BOISDUVAL et GUENÉE. Deltoites et Pyralites. Par M. A. GUENÉE. Paris, 448 pp., 10 pls.
- GUENÉE M. A. 1862. Annexe G de l'ouvrage intitulé: Notes sur l'Île de la Réunion par L. Maillard. Lépidoptères. G. 70.
- GUILDING L. 1828. Insects infesting the sugar cane. Trans. Soc. Arts, London, **46**: 43—153.
- HAIMBACH F. 1907. Two new species of *Crambus* and a new variety of *Haematopsis grataria* FABRICIUS. Ent. News, Philadelphia, **18**: 45—45.
- HAMPSON G. F. 1893. Illustrations of typical specimens of *Lepidoptera Heterocera* in the collection of the British Museum. Part. ix. The Macrolepidoptera Heterocera of Ceylon. London, 4to, V+182 pp., pls. clvii—clxxvi.
- HAMPSON G. F. 1896. On the classification of the *Schoenobiinae* and *Crambinae*, two Subfamilies of Moths, of the Family *Pyralidae*. Proc. zool. Soc. Lond., London, **1895**: 897—974.
- HAMPSON G. F. 1912. The Moths of India. Supplementary Paper to the volume in „The Fauna of British India“, Ser. iv. Part v. J. Nat. Hist., Bombay, **21**: 1222—1272, pl. G.
- HAMPSON G. F. 1919. Descriptions of new *Pyralidae* of the subfamilies *Crambinae* and *Siginae*. Ann. Mag. nat. Hist., London, (9) **3**: 275—292, 437—457, 533—547; (9) **4**: 53—58, 137—154, 305—326.
- HAWORTH A. H. 1811. *Lepidoptera Britannica*. (3)pp. 377-512. London.
- HEINEMANN H. 1865. Die Schmetterlinge Deutschlands und der Schweiz. Zweite Abtheilung. Kleinschmetterlinge. Band I. Heft II. Die Zünsler. Braunschweig, VI+214 pp.
- HERRICH-SCHÄFFER G. A. W. 1848—1849. Systematischer Bearbeitung der Schmetterlinge von Europa zugleich als Text, Revision und Supplement, zu Jakob HÜBNER's Sammlung europäischer Schmetterlinge. Vierter Band. Die Zünsler und Wickler. Regensburg, 288 pp.
- HERRICH-SCHÄFFER G. A. W. 1850—1858. Sammlung neuer oder wenig bekannter aussereuropäischer Schmetterlinge. Regensburg, 84 pp., 100 pls.
- HULST G. D. 1886. Descriptions of new *Pyralidae*. Trans. Am. ent. Soc., Washington, **13**: 145—168.
- HULST G. D. 1887. Notes upon certain *Pyralidae*. Ent. Amer., Philadelphia, **3**: 21—22.
- HULST G. D. 1888. New Genera and Species of *Epipaschia* and *Phycitidae*. Ent. Am., Philadelphia, **4**: 113—116.
- HULST G. D. 1900. New Species of *Lepidoptera*. J. N. York ent. Soc., New York, **8**: 215—225.
- HÜBNER J. 1796—[1838]. Sammlung europäischer Schmetterlinge. Horde viii. Tineae-Schaben. Augsburg, 78 pp., 69 pls., pls. lxx and lxxvi publ. by GEYER.

- HÜBNER J. 1816—[1826]. Verzeichniss bekannter Schmettlinge [sic]. Augsburg, 432 pp.
- HÜBNER J. 1818. Zuträge zur Sammlung exotischer Schmetterlinge bestehend in Bekundigung einzelner Fliegmuster neuer oder rarer nichteuropäischer Gattungen. Erstes Hundert. Augsburg, pp. 1—40, 1—172, pls. 492—663.
- HÜBNER J. [1819—1827]. Sammlung exotischer Schmetterlinge errichtet. Zweiter Band. Augsburg, 4 pp. of index, 225 col. pls.
- JOANNIS J. 1910. Description d'un nouveau genre de *Galleriinae* de France. Bull. Soc. ent. Fr., Paris, **1910**: 270—271.
- JOANNIS J. 1932. Lépidoptères hétérocères des Mascareignes. Soc. ent. Fr., Paris, Livre centen.: 427—456, 1 pl.
- JOANNIS L., RAGONOT E.-L. Description de Genres nouveaux et espèces nouvelles de Lépidoptères. Deuxieme partie par E.-L. RAGONOT. Ann. Soc. ent. Fr., Paris, (6) **8**: 275—284, pl. vi.
- KAPUR A. P. 1948. A note on the genus *Epina* WALKER (*Lep.*, *Pyral.*). Ent., London, **81**: 191.
- KAPUR A. P. 1950. The identity of some *Crambinae* associated with sugar-cane in India and of certain species related to them (*Lepidoptera*, *Pyralidae*). Trans. ent. Soc. Lond., London, **101**: 289—434, 10 pls.
- KEARFOTT W. D. 1908. Descriptions of new species of North American crambid moths. Proc. U. S. nat. Mus., Washington, **35**: 367—393.
- KIRBY W. F. 1897. A hand-book to the order Lepidoptera. Allen's Naturalist's Library, vol. v. Moths. Part iii. London, xii+332 pp., pls. cxvii—clviii.
- LATTIN G. de. 1952. Studien über die Gattung *Crambus* F. ii. Über die Gattungszugehörigkeit des „*Crambus*“ *malacellus* DUR. Ent. Zeit., Stuttgart, **62**: 89, 91.
- LATTIN G. de. 1961. Eine Notiz zur Gattungs-Nomenklatur der Crambiden. Ent. Zeit., Stuttgart, **71**: 115.
- LEECH J. H. 1889. New species of Crambi from Japan and Corea. Ent., London, **22**: 106—109, pl. v.
- LHOMME L. 1935. Catalogue des Lépidoptères de France et de Belgique. Vol. 2, fasc. 1. Le Carriol, 172 pp.
- LINNAEUS C. 1758. *Systema Naturae...* Edition decima, reformata. Tomus I, *Regnum Animale*. Holmiae, ii+824 pp.
- MABILLE P. 1899. *Lepidoptera nova Malgassica et Africana*. Ann. Soc. ent. Fr., Paris, **68**: 723—753.
- MABILLE P. 1906. Notes sur plusieurs Lépidoptères de la faune paléarctique. Ann. Soc. ent. Fr., Paris, **75**: 31—36.
- MARION H. 1954. Contribution à l'étude des *Pyralidae* de Madagascar. Mém. Inst. sci. Madag., Tananarive, (E) **5**: 39—62, 1 pl.
- MARTIN E. L. 1956. On the Systematic Position and Status of Some Pyralid Genera and Species. Ent., London, **89**: 163—166.
- MARUMO N. 1933. Studies on Rice Borers. I. Classification of the subfamily *Crambinae* in Japan. Tokyo, 61 pp., 6 pls, 4 pp.
- MATSUMURA F. 1911. Erster Beitrag zur Insecten-Fauna Sachalin. J. Coll. Agric. Hokkaido Imp. Univ. Sapporo, **4**: 1—145, 2 pls.
- MATSUMURA F. 1925. An enumeration of the butterflies and moths from Saghalien, with descriptions of new species and subspecies. J. Coll. Agric. Hokkaido Imp. Univ., Sapporo, **15**: 83—196, 4 pls.
- MEYRICK E. 1879. Descriptions of Australian *Micro-Lepidoptera*: II. *Crambites*. Proc. Linn. Soc. N. S. Wales, Sydney, **4**: 205—242.
- MEYRICK E. 1882. Descriptions of New Zealand *Micro-Lepidoptera*. (I.). N. Zeal. J. Sc., Dunedin, **1**: 186—187.
- MEYRICK E. 1882. Descriptions of Australian *Microlepidoptera*. Proc. linn. Soc. N. S. Wales, Sydney, **7**: 148—202.

- MEYRICK E. 1884. On the classification of the Australian *Pyralidina*. Trans. ent. Soc. Lond., London, **1884**: 277—350.
- MEYRICK E. 1885. Descriptions of New Zealand *Micro-Lepidoptera*. Trans. N. Zeal. Inst., Wellington, **17**: 68—149.
- MEYRICK E. 1886. Descriptions of *Lepidoptera* from the South Pacific. Trans. ent. Soc. Lond., London, **1886**: 189—296.
- MEYRICK E. 1887. Descriptions of some exotic *Micro-Lepidoptera*. Trans. ent. Soc. Lond., London, **1887**: 269—280.
- MEYRICK E. 1888. Notes on New Zealand *Pyralidina*. Trans. N. Zeal. Inst., Wellington, **20**: 62—73.
- MEYRICK E. 1889. On some *Lepidoptera* from New Guinea. Trans. ent. Soc. Lond., London, **1889**: 455—522.
- MEYRICK E. 1897a. Descriptions of new *Lepidoptera* from Australia and New Zealand. Trans. ent. Soc. Lond., London, **1897**: 367—390.
- MEYRICK E. 1897b. On *Lepidoptera* from the Malay Archipelago. Trans. ent. Soc. Lond., London, **1897**: 69—92.
- MEYRICK E. 1899. *Macrolepidoptera*. In: Fauna Hawaïensis. **1(2)**: 123—275, pls. iii—vii, Cambridge.
- MEYRICK E. 1909. *Lepidoptera* from Auckland Island. Subantarct. Isl. New Zealand, **1**: 70—74.
- MEYRICK E. 1930—1937. Exotic *Microlepidoptera*. **4**: 1—32 (1930); 33—192 (1931); 193—252 (1932); 253—448 (1933); 449—544 (1934); 545—608 (1935); 609—942 + index (1936); **5**: 1—64 (1936); 65—160 + index (1937). London.
- MEYRICK E. 1936. New species of Pyrales and Microlepidoptera from the Deutsches Entomologisches Institut. Arb. morph. taxon. Ent., Berlin-Dahlem, **3**: 94—109.
- MOORE F. 1866. The *Lepidoptera* of Ceylon. London, **3**: 305—392, pls. clxxxii—excv.
- MORRISON H. K. 1874. Descriptions of New *Noctuidae*. Proc. Bost. Soc. nat. Hist., Boston, **17**: 131—166.
- MÖSCHLER H. B. 1883. Beiträge zur Schmetterlings-Fauna von Surinam. V. (Supplement). Verh. zool.-bot. Ges., Wien, **32**: 303—362, pls. xvii—xviii.
- MÖSCHLER H. B. 1890. Die Lepidopteren-Fauna von Portorico. Abh. Senck. Ges., Frankfurt, **16**: 69—360, 1 pl.
- OKANO M. 1958. Three New Species of the Japanese *Crambinae* (*Lepidoptera*, *Pyralididae*). *Tinea*, **4**: 259—262, 6 Figs.
- OKANO M. 1962a. Notes on Some Japanese *Crambinae* (*Pyralididae*) (4). Trans Lep. Soc. Japan, Tokyo, **12**: 51.
- OKANO M. 1962b. The systematic study of the Japanese *Crambinae*. Ann. Lep. Gak. Fac. Iwate Univ., **20**: 83—137, 15 pls.
- PHILPOTT A. 1930. Descriptions of Lepidoptera in the Canterbury Museum. Rec. Canterbury Mus., Christchurch, N. Z., **3**: 247—250.
- PURDIE A. 1884. *Orocrambus*, sp. N. Zeal. J. Sci., Dunedin, **2**: 167—168.
- RAGONOT E. L., see JOANNIS L., RAGONOT E. L.
- RAGONOT E. L. 1892. In STAUDINGER: Lepidopteren aus Tunis. Dtsch. ent. Zeit. Lep., Berlin, **5**: 277—300.
- REBEL H. 1892. Beitrag zur Microlepidopteren-fauna des canarischen Archipels. Ann. Hofmus. Wien, Wien, **7**: 241—284, pl. xvii.
- ROBINSON C. T. 1871. Lepidopterological Miscellanies No. 2. Ann. Lyc. N. York, New York, **9**: 310—316.
- ROSENSTOCK R. 1885. Notes on Australian *Lepidoptera*, with descriptions of new Species. Ann. Mag. Nat. Hist., London, (5) **16**: 421—443, pl. xi.
- SALMON J. T., BRADLEY J. D. 1956. *Lepidoptera* from the Cape Expedition and Antipodes Islands. Rec. Dom. Mus., Wellington, **3** (1): 61—81.
- SCHAUS W. 1913. New species of *Heterocera* from Costa Rica. xix. Ann. Mag. nat. Hist., London, (8) **11**: 1—43, 234—262, 342—358, 361—386.

- SCHAUS W. 1922. New Species of *Pyralidae* of the Subfamily *Crambinae* from tropical America. Proc. ent. Soc. Wash., Washington D. C., **24**: 127—145.
- SCHAUS W. 1924. New species of moths in the United States National Museum. Proc. U. S. nat. Mus., Washington D. C., **65** (7), 74 pp.
- SCHAWERDA C. 1937. Zwei neue Microheteroceren aus Mazedonien. Zeit. öst. ent. Ver., Wien, **22**: 55—56.
- SCOPOLI I. A. 1763. Entomologia Carniolica exhibens Insecta Carnioliae indigena et distributa in ordines, genera, species, varietates. Methodo Linnaeana. Vindobonae, 421 pp.
- SHIBUYA J. 1928. I. On the Japanese *Crambinae*. II. The systematic study on the Japanese *Pyralinae*. (Lepid.). J. Fac. Agr., Sapporo, **21**: 121—176, 3 pls.
- SNELLEN P. C. T. 1880. Nieuwe Pyraliden op het eiland Celebes gevonden. Tijd. Ent., **23**: 198—250.
- SNELLEN P. C. T. 1890a. A catalogue of the *Pyralidina* of Sikkim collected by Henry J. ELWES and the late Otto MöLLER. Trans. ent. Soc. Lond., London, **1890**: 557—647, pls. xix, xx.
- SNELLEN P. C. T. 1890b. Meded. Proefst. „Kagok“. **1890**: 94.
- SNELLEN P. C. T. 1895. Anteekenigen over Pyraliden beschrijving van nieuwe genera en soorten. Tijd. Ent., Amsterdam, **38**: 103—161.
- STEPHENS J. F. 1834. Illustrations of British Entomology or, a Synopsis of Indigenous Insects: containing their generic and specific distinctions, vol. IV. London, 436 pp., pls. xxiii—xl.
- STRAND E. 1907. Rectification synonymique. Bull. Soc. ent. Fr., Paris, **1907**: 175.
- STRAND E. 1919. H. SAUTER'S Formosa-Ausbeute: *Pyralidae*, Subfam. *Gallerinae*, *Crambinae*, *Schoenobiinae*, *Anerastiinae* und *Phycitiinae*. Ent. Ztg., Stettin, **79**: 248—276.
- SWINHOE C. 1885. On the Lepidoptera of Bombay and the Deccan. Part. IV. *Heterocera*, continued. Proc. zool. Soc. Lond. **1885**: 852—886, pls. lvi—lvii.
- TAMS W. H. T. 1942. Note on the name of the sugar-cane borer of Mauritius (*Lep.*, *Pyral.*). Bull. ent. Res., London, **33**: 67—68.
- TURNER A. J. 1904. A preliminary revision of the Australian *Thyrididae* and *Pyralidae*. Part. I. Proc. roy. Soc. Queensland, Brisbane, **18**: 160—199.
- TURNER A. J. 1905. A preliminary revision of the Australian *Thyrididae* and *Pyralidae*. Part. II. Proc. roy. Soc. Queensland, Brisbane, **19**: 39—63.
- TURNER A. J. 1911. Studies in Australian Lepidoptera. Ann. Queensland Mus., Brisbane, **10**: 59—135.
- TURNER A. J. 1922. Studies in Australian Lepidoptera. Proc. roy. Soc. Victoria, Melbourne, **35** (N. S.): 26—62.
- TURNER A. J. 1932. New Australian Lepidoptera. Trans. roy. Soc. South Austr., **56**: 175, 196.
- TURNER A. J. 1937. New Australian *Pyraloidea* (Lepidoptera). Proc. roy. Soc. Queensland, Brisbane, **48**: 61—88.
- WALKER F. 1856—1866. List of the Specimens of Lepidopterous Insects in the Collection of the British Museum. London. Part IX (1856), 252 pp; part XVII (1859), pp. 255—508; part XXVII (1863) pp. 1—286; part XXX (1864) pp. 837—1096; part XXXV (1866), pp. 1545—2030.
- WALLENGREN H. D. J. 1868. *Lepidoptera*. Species novas descriptis. In: Konglja Svenska Fre-gatten Eugenies resa omkring Jorden under befäl of C. A. VIRGIN. Aren 1851—1853. Andra delen Zoologi. 1. *Insecta*. Stockholm, pp. 351—390, pls. vi, vii.
- WARREN W. 1891—1892. Descriptions of new Genera and Species of *Pyralidae* contained in the British Museum Collection. Ann. Mag. nat. Hist., London, (6) **8**: 61—70; (6) **9**: 172.
- WOLLASTON T. V. 1879. Notes on the Lepidoptera of St. Helena. Ann. Mag. nat. Hist., London, (5) **3**: 219—233, 329—343, 415—441.
- VIETTE P. 1960. Pyrales de Madagascar et des Comores nouvelles ou peu connues. Rev. fr. d'Ent., Paris, **27** (2): 200—214.
- ZELLER P. C. 1839. Versuch einer naturgemassen Eintheilung der Schaben. Isis v. Oken, Jena-Leipzig, **1839**: 167—219.
- ZELLER P. C. 1846. Die Arten der Gattung *Eudorea*. Lnn. Ent., Berlin, **1**: 262—318.

- ZELLER P. C. 1847. Bemerkungen über die auf einer Reise nach Italien und Sicilien beobachteten Schmetterlingsarten. *Isis von Oken*, Jena-Leipzig, **1847**: 721—771.
- ZELLER P. C. 1849. Verzeichniss der von Herrn Jos. MANN beobachteten Toscanischen Microlepidopteren. *Ent. Ztg.*, Stettin, **10**: 312—317.
- ZELLER P. C. 1863. *Chilonidarum et Crambidarum genera et species*. Berolinensis, 56 pp.
- ZELLER P. C. 1866. Beschreibung einiger amerikanischen Wickler und Crambiden. *Ent. Ztg.*, Stettin, **27**: 137—157, 1 pl. i.
- ZELLER P. C. 1872. Beiträge zur Kenntniss der nordamerikanischen Nachtfalter besonders der Microlepidopteren. *Verh. zool.-bot. Ges.*, Wien, **22**: 447—566, pls. ii, iii.
- ZELLER P. C. 1877. Exotische Microlepidopteren. *Horae Soc. ent. ross.*, St. Petersbourg, **13**: 3—493, pls. i—vi.
- ZERNY H. 1914. Über paläarktische Pyraliden des k. k. naturhistorischen Hofmuseum in Wien. *Ann. K. K. nat.-hist. Mus.*, Wien, **28**: 295—348, pls. xxv & xxvi.
- ZIMMERMAN E. C. 1958. Insects of Hawaii. A Manual of the Insects of the Hawaiian Islands, including an Enumeration of the Species and Notes on their Origin, Distribution, Hosts, Parasites, etc. Vol. 8. *Lepidoptera: Pyraloidea*. Honolulu, viii + 456 pp.
- ZINCKEN J. L. T. F. 1817. Die Linneischen Tineen in ihre natürlichen Gattungen aufgelöst und beschrieben. *GERMAR's Magazin der Entomologie*. Zweiter Band : 24—113, Halae.

STRESZCZENIE

Autor przedstawia wykaz rodzajów *Crambidae*, podając dane odnośnie do ich synonimiki i typów rodzajowych.

РЕЗЮМЕ

Автор дает перечень родов *Crambidae* приводя данные касающиеся их синонимики и генеративных типов.

Redaktor zeszytu: dr W. Szymczakowski

Państwowe Wydawnictwo Naukowe — Oddział w Krakowie 1963

Nakład 800+100 egz. — Ark. wyd. 3 — Ark. druk. 2¹⁰/₁₆ — Papier ilustr. kl. III 80 g 70×100
Zam. 631/62 Cena zł 12,—

Drukarnia Uniwersytetu Jagiellońskiego w Krakowie